arXiv:1106.1513v1 [math.FA] 8 Jun 2011

DILATION OF RITT OPERATORS ON L^p-SPACES

CÉDRIC ARHANCET, CHRISTIAN LE MERDY

ABSTRACT. For any Ritt operator $T\colon L^p(\Omega) \to L^p(\Omega)$, for any positive real number α, and for any $x \in L^p(\Omega)$, we consider $\|x\|_{T,\alpha} = \Big\| \Big(\sum_{k=1}^{\infty} k^{2\alpha-1} |T^{k-1}(I-T)^\alpha x|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p}$. We show that if T is actually an R-Ritt operator, then the square functions $\| \ \|_{T,\alpha}$ are pairwise equivalent. Then we show that T and its adjoint $T^*\colon L^{p'}(\Omega) \to L^{p'}(\Omega)$ both satisfy uniform estimates $\|x\|_{T,1} \lesssim \|x\|_{L^p}$ and $\|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}}$ for $x \in L^p(\Omega)$ and $y \in L^{p'}(\Omega)$ if and only if T is R-Ritt and admits a dilation in the following sense: there exist a measure space $\widetilde{\Omega}$, an isomorphism $U\colon L^p(\widetilde{\Omega}) \to L^p(\widetilde{\Omega})$ such that $\{U^n : n \in \mathbb{Z}\}$ is bounded, as well as two bounded maps $L^p(\Omega) \xrightarrow{J} L^p(\widetilde{\Omega}) \xrightarrow{Q} L^p(\Omega)$ such that $T^n = QU^nJ$ for any $n \geq 0$. We also investigate functional calculus properties of Ritt operators and analogs of the above results on noncommutative L^p-spaces.

2000 Mathematics Subject Classification : 47B38, 47A20, 47A60.

1. INTRODUCTION.

Let (Ω, μ) be a measure space and let $1 < p < \infty$. For any bounded operator $T\colon L^p(\Omega) \to L^p(\Omega)$, consider the 'square function'

$$\|x\|_{T,1} = \Big\| \Big(\sum_{k=1}^{\infty} k \big| T^k(x) - T^{k-1}(x) \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p}, \tag{1.1}$$

defined for any $x \in L^p(\Omega)$. Such quantities frequently appear in the analysis of L^p-operators. They go back at least to [50], where they were used in connection with martingale square functions to study diffusion semigroups and their discrete counterparts. Similar square functions for continuous semigroups played a key role in the recent development of H^∞-calculus and its applications. See in particular the fundamental paper [12], the survey [31] and the references therein.

It is shown in [34] that if T is both a positive contraction and a Ritt operator, then it satisfies a uniform estimate $\|x\|_{T,1} \lesssim \|x\|_{L^p}$ for $x \in L^p(\Omega)$. This estimate and related ones lead to strong maximal inequalities for this class of operators (see also [35]). Next in the paper [30], the second named author studies the operators T such that both $T\colon L^p(\Omega) \to L^p(\Omega)$ and its adjoint operator $T^*\colon L^{p'}(\Omega) \to L^{p'}(\Omega)$ satisfy uniform estimates

$$\|x\|_{T,1} \lesssim \|x\|_{L^p} \qquad \text{and} \qquad \|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}} \tag{1.2}$$

Date: January 28, 2013.

for $x \in L^p(\Omega)$ and $y \in L^{p'}(\Omega)$. (Here $p' = \frac{p}{p-1}$ is the conjugate number of p.) It is shown that (1.2) implies that T is an R-Ritt operator (see Section 2 below for the definition) and that (1.2) is equivalent to T having a bounded H^∞-calculus with respect to a Stolz domain of the unit disc with vertex at 1.

The present paper is a continuation of these investigations. Our main result is a characterization of (1.2) in terms of dilations. We show that (1.2) holds true if and only if T is R-Ritt and there exist another measure space $(\widetilde{\Omega}, \widetilde{\mu})$, two bounded maps $J\colon L^p(\Omega) \to L^p(\widetilde{\Omega})$ and $Q\colon L^p(\widetilde{\Omega}) \to L^p(\Omega)$, as well as an isomorphism $U\colon L^p(\widetilde{\Omega}) \to L^p(\widetilde{\Omega})$ such that $\big\{U^n \,:\, n \in \mathbb{Z}\big\}$ is bounded and

$$T^n = QU^nJ, \qquad n \geq 0.$$

This result will be established in Section 4. It should be regarded as a discrete analog of the main result of [19].

In Section 3, we consider variants of (1.1) as follows. Assume that $T\colon L^p(\Omega) \to L^p(\Omega)$ is a Ritt operator. Then $I - T$ is a sectorial operator and one can define its fractional power $(I-T)^\alpha$ for any $\alpha > 0$. Then we consider

$$\|x\|_{T,\alpha} \,=\, \bigg\| \Big(\sum_{k=1}^\infty k^{2\alpha-1} \big| T^{k-1}(I-T)^\alpha x \big|^2 \Big)^{\frac{1}{2}} \bigg\|_{L^p} \tag{1.3}$$

for any $x \in L^p(\Omega)$. Our second main result (Theorem 3.3 below) is that when T is an R-Ritt operator, then the square functions $\| \ \|_{T,\alpha}$ are pairwise equivalent. This result of independent interest should be regarded as a discrete analog of [33, Thm. 1.1]. We prove it here as it is a key step in our characterization of (1.2) in terms of dilations.

Section 2 mostly contains preliminary results. Section 5 is devoted to complements on L^p-operators and their functional calculus properties, in connection with p-completely bounded maps. Finally Section 6 contains generalizations to operators $T\colon X \to X$ on general Banach spaces X. We pay a special attention to noncommutative L^p-spaces, in the spirit of [22].

We end this introduction with a few notation. If X is a Banach space, we let $B(X)$ denote the algebra of all bounded operators on X and we let I_X denote the identity operator on X (or simply I if there is no ambiguity on X). For any $T \in B(X)$, we let $\sigma(T)$ denote the spectrum of T. If $\lambda \in \mathbb{C} \setminus \sigma(T)$ (the resolvent set of T), we let $R(\lambda, T) = (\lambda I_X - T)^{-1}$ denote the corresponding resolvent operator. We refer the reader to [14] for general information on Banach space geometry. We will frequently use Bochner spaces $L^p(\Omega; X)$, for which we refer to [15].

For any $a \in \mathbb{C}$ and $r > 0$, we let $D(a,r) = \{z \in \mathbb{C} \,:\, |z-a| < r\}$ and we let $\mathbb{D} = D(0,1)$ denote the open unit disc centered at 0. Also we let $\mathbb{T} = \{z \in \mathbb{C} \,:\, |z| = 1\}$ denote its boundary.

Whenever $\mathcal{O} \subset \mathbb{C}$ is a non empty open set, we let $H^\infty(\mathcal{O})$ denote the space of all bounded holomorphic functions $f\colon \mathcal{O} \to \mathbb{C}$. This is a Banach algebra for the norm

$$\|f\|_{H^\infty(\mathcal{O})} \,=\, \sup\big\{ |f(z)| \,:\, z \in \mathcal{O} \big\}.$$

Also we let $\mathcal{P}$ denote the algebra of all complex polynomials.

In the above presentation and later on in the paper we will use $\lesssim$ to indicate an inequality up to a constant which does not depend on the particular element to which it applies. Then $A(x) \approx B(x)$ will mean that we both have $A(x) \lesssim B(x)$ and $B(x) \lesssim A(x)$.

2. Preliminaries on R-boundedness and Ritt operators.

This section is devoted to definitions and preliminary results involving R-boundedness (and the companion notion of γ-boundedness), matrix estimates and Ritt operators. We deal with operators acting on an arbitrary Banach space X (as opposed to the next two sections, where X will be an L^p-space).

Let $(\varepsilon_k)_{k\geq 1}$ be a sequence of independent Rademacher variables on some probability space Ω_0. We let $\mathrm{Rad}(X) \subset L^2(\Omega_0; X)$ be the closure of $\mathrm{Span}\big\{\varepsilon_k \otimes x \ : \ k \geq 1, \ x \in X\big\}$ in the Bochner space $L^2(\Omega_0; X)$. Thus for any finite family $x_1, \ldots, x_n$ in X, we have

$$\Big\| \sum_{k=1}^n \varepsilon_k \otimes x_k \Big\|_{\mathrm{Rad}(X)} = \Big(\int_{\Omega_0} \Big\| \sum_{k=1}^n \varepsilon_k(\omega)\, x_k \Big\|_X^2 \, d\omega \Big)^{\frac{1}{2}}.$$

We say that a set $F \subset B(X)$ is R-bounded provided that there is a constant $C \geq 0$ such that for any finite families $T_1, \ldots, T_n$ in F and $x_1, \ldots, x_n$ in X, we have

$$\Big\| \sum_{k=1}^n \varepsilon_k \otimes T_k(x_k) \Big\|_{\mathrm{Rad}(X)} \leq \ C \, \Big\| \sum_{k=1}^n \varepsilon_k \otimes x_k \Big\|_{\mathrm{Rad}(X)}.$$

In this case we let $R(F)$ denote the smallest possible C, which is called the R-bound of F.

Let $(g_k)_{k\geq 1}$ denote a sequence of independent complex valued, standard Gaussian random variables on some probability space Ω_1, and let $\mathrm{Gauss}(X) \subset L^2(\Omega_1; X)$ be the closure of $\mathrm{Span}\big\{g_k \otimes x \ : \ k \geq 1, \ x \in X\big\}$. Then replacing the ε_k's and $\mathrm{Rad}(X)$ by the g_k's and $\mathrm{Gauss}(X)$ in the above paragraph, we obtain the similar notion of γ-bounded set. The corresponding γ-bound of a set F is denoted by $\gamma(F)$.

These two notions are very close to each other, however we need to work with both of them in this paper. Comparing them, we recall that any R-bounded set $F \subset B(X)$ is automatically γ-bounded, with $\gamma(F) \leq R(F)$. Moreover if X has a finite cotype, then the Rademacher averages and the Gaussian averages are equivalent on X (see e.g. [14, Prop. 12.11 and Thm. 12.27]), hence F is R-bounded if (and only if) it is γ-bounded.

R-boundedness was introduced in [5] and then developed in the fundamental paper [9]. We refer to the latter paper and to [26, Section 2] for a detailed presentation. We recall two facts which are highly relevant for our paper. First, the closure of the absolute convex hull of any R-bounded set is R-bounded [9, Lem. 3.2]. This implies the following.

Lemma 2.1. *Let $F \subset B(X)$ be an R-bounded set, let $J \subset \mathbb{R}$ be an interval and let $C \geq 0$ be a constant. Then the set*

$$\Big\{ \int_J a(t) V(t)\, dt \ \Big| \ V\colon J \to F \text{ is continuous, } a \in L^1(J), \ \|a\|_{L^1(J)} \leq C \Big\}$$

is R-bounded.

Second, if $X = L^p(\Omega)$ is an L^p-space with $1 \le p < \infty$, then X has a finite cotype and we have an equivalence

$$\Big\| \sum_k \varepsilon_k \otimes x_k \Big\|_{\mathrm{Rad}(L^p(\Omega))} \approx \Big\| \Big(\sum_k |x_k|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p(\Omega)} \tag{2.1}$$

for finite families $(x_k)_k$ of $L^p(\Omega)$. Consequently a set $F \subset B\big(L^p(\Omega)\big)$ is R-bounded if and only if it is γ-bounded, if and only if there exists a constant $C \ge 0$ such that for any finite families $T_1, \ldots, T_n$ in F and $x_1, \ldots, x_n$ in $L^p(\Omega)$, we have

$$\Big\| \Big(\sum_{k=1}^n |T_k(x_k)|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p(\Omega)} \le C \, \Big\| \Big(\sum_{k=1}^n |x_k|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p(\Omega)} .$$

In the sequel we represent any element of $B(\ell^2)$ by an infinite matrix $[c_{ij}]_{i,j\ge 1}$ in the usual way. Likewise for any integer $n \ge 1$, we identify the algebra M_n of all $n \times n$ matrices with the space of linear maps $\ell_n^2 \to \ell_n^2$. Clearly an infinite matrix $[c_{ij}]_{i,j\ge 1}$ represents an element of $B(\ell^2)$ (in the sense that it is the matrix associated to a bounded operator $\ell^2 \to \ell^2$) if and only if

$$\sup_{n \ge 1} \big\| [c_{ij}]_{1 \le i,j \le n} \big\|_{B(\ell_n^2)} < \infty .$$

For any $[c_{ij}]_{1\le i,j\le n}$ in M_n, we set

$$\big\| [c_{ij}] \big\|_{\mathrm{reg}} = \big\| \big[|c_{ij}| \big] \big\|_{B(\ell_n^2)}.$$

This is the so-called 'regular norm' of the operator $[c_{ij}] \colon \ell_n^2 \to \ell_n^2$.

Lemma 2.2. *For any matrix $[c_{ij}]$ in M_n, the following assertions are equivalent.*

(i) *We have $\big\| [c_{ij}] \big\|_{\mathrm{reg}} \le 1$.*

(ii) *There exist two matrices $[a_{ij}]$ and $[b_{ij}]$ in M_n such that $c_{ij} = a_{ij} b_{ij}$ for any $i, j = 1, \ldots, n$, and we both have*

$$\sup_{1\le i\le n} \sum_{j=1}^n |a_{ij}|^2 \le 1 \qquad \text{and} \qquad \sup_{1\le j\le n} \sum_{i=1}^n |b_{ij}|^2 \le 1.$$

The implication '(ii)⇒(i)' is an easy application of the Cauchy-Schwarz inequality. The converse is due to Peller [42, Section 3] (see also [1]). We refer to [44] and [48, Sect. 1.4] for more about this result and complements on regular norms.

The following result extends the boundedness of the Hilbert matrix (wich corresponds to the case $\beta = \gamma = \frac{1}{2}$). We thank Éric Ricard for his precious help in devising this proof.

Proposition 2.3. *Let $\beta, \gamma > 0$ be two positive real numbers. Then the infinite matrix*

$$\left[\frac{i^{\beta - \frac{1}{2}} j^{\gamma - \frac{1}{2}}}{(i+j)^{\beta+\gamma}} \right]_{i,j\ge 1}$$

represents an element of $B(\ell^2)$.

Proof. For any $i, j \geq 1$, set

$$c_{ij} = \frac{i^{\beta-\frac{1}{2}} j^{\gamma-\frac{1}{2}}}{(i+j)^{\beta+\gamma}}, \qquad a_{ij} = c_{ij}^{\frac{1}{2}} \big(\tfrac{i}{j}\big)^{\frac{1}{4}}, \qquad \text{and} \qquad b_{ij} = c_{ij}^{\frac{1}{2}} \big(\tfrac{j}{i}\big)^{\frac{1}{4}}.$$

Then $c_{ij} = a_{ij} b_{ij}$ for any $i, j \geq 1$, hence by the easy implication of Lemma 2.2, it suffices to show that

$$\sup_{i \geq 1} \sum_{j=1}^{\infty} |a_{ij}|^2 < \infty \qquad \text{and} \qquad \sup_{j \geq 1} \sum_{i=1}^{\infty} |b_{ij}|^2 < \infty. \tag{2.2}$$

Fix some $i \geq 1$. For any $j \geq 1$, we have

$$|a_{ij}|^2 = c_{ij} \big(\tfrac{i}{j}\big)^{\frac{1}{2}} = \frac{i^{\beta} j^{\gamma-1}}{(i+j)^{\beta+\gamma}}.$$

Hence

$$\sum_{j=1}^{\infty} |a_{ij}|^2 = i^{\beta} \left(\frac{1}{(i+1)^{\beta+\gamma}} + \sum_{j=2}^{\infty} \frac{1}{j^{1-\gamma}(i+j)^{\beta+\gamma}} \right).$$

Looking at the variations of the function $t \mapsto 1/(t^{1-\gamma}(i+t)^{\beta+\gamma})$ on $(1, \infty)$, we immediately deduce that

$$\sum_{j=1}^{\infty} |a_{ij}|^2 \leq 1 + 2\, i^{\beta} \int_1^{\infty} \frac{1}{t^{1-\gamma}(i+t)^{\beta+\gamma}}\, dt.$$

Changing t into it in the latter integral, we deduce that

$$\sum_{j=1}^{\infty} |a_{ij}|^2 \leq 1 + 2 \int_0^{\infty} \frac{1}{t^{1-\gamma}(1+t)^{\beta+\gamma}}\, dt.$$

This upper bound is finite and does not depend on i, which proves the first half of (2.2). The proof of the second half is identical. □

We record the following elementary lemma for later use.

Lemma 2.4. *Let $[c_{ij}]_{i,j \geq 1}$ and $[d_{ij}]_{i,j \geq 1}$ be infinite matrices of nonnegative real numbers, such that $c_{ij} \leq d_{ij}$ for any $i, j \geq 1$. If the matrix $[d_{ij}]_{i,j \geq 1}$ represents an element of $B(\ell^2)$, then the same holds for $[c_{ij}]_{i,j \geq 1}$.*

We will need the following classical fact (see e.g. [14, Cor. 12.17]).

Lemma 2.5. *Let X be a Banach space and let $[b_{ij}]_{1 \leq i,j \leq n}$ be an element of M_n. Then for any $x_1, \ldots, x_n$ in X, we have*

$$\left\| \sum_{i,j=1}^{n} g_i \otimes b_{ij} x_j \right\|_{\mathrm{Gauss}(X)} \leq \big\| [b_{ij}] \big\|_{B(\ell_n^2)} \left\| \sum_{j=1}^{n} g_j \otimes x_j \right\|_{\mathrm{Gauss}(X)}.$$

That result does not remain true if we replace Gaussian variables by Rademacher variables and this defect is the main reason why it is sometimes easier to deal with γ-boundedness than with R-boundedness.

Proposition 2.6. *Let X be a Banach space, let $F = \{T_{ij} : i,j \geq 1\}$ be a γ-bounded family of operators on X, let $n \geq 1$ be an integer and let $[c_{ij}]_{1\leq i,j\leq n}$ be an element of M_n. Then for any $x_1, \ldots, x_n$ in X, we have*

$$\Big\| \sum_{i,j=1}^n g_i \otimes c_{ij} T_{ij}(x_j) \Big\|_{\mathrm{Gauss}(X)} \leq \gamma(F) \, \big\| [c_{ij}] \big\|_{\mathrm{reg}} \Big\| \sum_{j=1}^n g_j \otimes x_j \Big\|_{\mathrm{Gauss}(X)}.$$

Proof. We can assume that $\big\| [c_{ij}] \big\|_{\mathrm{reg}} \leq 1$. By Lemma 2.2, we can write $c_{ij} = a_{ij} b_{ij}$ with

$$\sup_{1\leq i\leq n} \sum_{j=1}^n |a_{ij}|^2 \leq 1 \qquad \text{and} \qquad \sup_{1\leq j\leq n} \sum_{i=1}^n |b_{ij}|^2 \leq 1. \tag{2.3}$$

Let $(g_{i,j})_{i,j\geq 1}$ be a doubly indexed family of independent Gaussian variables. For any integers $1 \leq i,j \leq n$, we define

$$A(i) = \begin{bmatrix} a_{i1} & a_{i2} & \ldots & a_{in} \end{bmatrix} \qquad \text{and} \qquad B(j) = \begin{bmatrix} b_{1j} & b_{2j} & \ldots & b_{nj} \end{bmatrix}^T.$$

Then we consider the two matrices

$$A = \mathrm{Diag}\big(A(1), \ldots, A(n)\big) \in M_{n,n^2} \qquad \text{and} \qquad B = \mathrm{Diag}\big(B(1), \ldots, B(n)\big) \in M_{n^2,n}.$$

Let $x_1, \ldots, x_n \in X$. Applying Lemma 2.5 successively to A and B, we then have

$$\begin{aligned} \Big\| \sum_{i,j=1}^n g_i \otimes c_{ij} T_{ij}(x_j) \Big\|_{\mathrm{Gauss}(X)} &= \Big\| \sum_{i,j=1}^n g_i \otimes a_{ij} b_{ij} T_{ij}(x_j) \Big\|_{\mathrm{Gauss}(X)} \\ &\leq \|A\| \, \Big\| \sum_{i,j=1}^n g_{ij} \otimes b_{ij} T_{ij}(x_j) \Big\|_{\mathrm{Gauss}(X)} \\ &\leq \gamma(F) \, \|A\| \, \Big\| \sum_{i,j=1}^n g_{ij} \otimes b_{ij} x_j \Big\|_{\mathrm{Gauss}(X)} \\ &\leq \gamma(F) \, \|A\| \|B\| \, \Big\| \sum_{j=1}^n g_j \otimes x_j \Big\|_{\mathrm{Gauss}(X)}. \end{aligned}$$

We have

$$\|A\| = \sup_{1\leq i\leq n} \big\| A(i) \big\|_{M_{1,n}} = \sup_{1\leq i\leq n} \Big(\sum_{j=1}^n |a_{ij}|^2 \Big)^{\frac{1}{2}},$$

hence $\|A\| \leq 1$ by (2.3). Likewise, we have $\|B\| \leq 1$ hence the above inequality yields the result. □

We now turn to Ritt operators, the key class of this paper, and recall some of their main features. Details and complements can be found in [6, 7, 30, 36, 38, 39, 52]. We say that an operator $T \in B(X)$ is a Ritt operator if the two sets

$$\{T^n : n \geq 0\} \qquad \text{and} \qquad \big\{ n(T^n - T^{n-1}) : n \geq 1 \big\} \tag{2.4}$$

are bounded. This is equivalent to the spectral inclusion

$$\sigma(T) \subset \overline{\mathbb{D}} \tag{2.5}$$

and the boundedness of the set

$$\big\{(\lambda-1)R(\lambda,T) \,:\, |\lambda|>1\big\}. \tag{2.6}$$

This resolvent estimate outside the unit disc is called the 'Ritt condition'.

Likewise we say that T is an R-Ritt operator if the two sets in (2.4) are R-bounded. This is equivalent to the inclusion (2.5) and the R-boundedness of the set (2.6).

For any angle $\gamma \in \big(0, \frac{\pi}{2}\big)$, let B_γ be the interior of the convex hull of 1 and the disc $D(0, \sin\gamma)$ (see Figure 1 below).



FIGURE 1.

Then the Ritt condition and its R-bounded version can be strengthened as follows.

Lemma 2.7. *Let $T\colon X \to X$ be a Ritt operator (resp. an R-Ritt operator). There exists an angle $\gamma \in (0, \frac{\pi}{2})$ such that*

$$\sigma(T) \subset B_\gamma \cup \{1\} \tag{2.7}$$

and the set

$$\big\{(\lambda-1)R(\lambda,T) \,:\, \lambda \in \mathbb{C}\setminus B_\gamma,\ \lambda \neq 1\big\} \tag{2.8}$$

is bounded (resp. R-bounded).

This essentially goes back to [6], see [30] for details.

For any angle $\theta \in (0,\pi)$, let

$$\Sigma_\theta = \big\{z \in \mathbb{C} \,:\, \big|\mathrm{Arg}(z)\big| < \theta\big\} \tag{2.9}$$

be the open sector of angle 2θ around the positive real axis $(0,\infty)$. We say that a closed operator $A\colon D(A) \to X$ with dense domain $D(A)$ is sectorial if there exists $\theta \in (0,\pi)$ such that $\sigma(A) \subset \overline{\Sigma_\theta}$ and the set

$$\{zR(z,A) \,:\, z \in \mathbb{C}\setminus \overline{\Sigma_\theta}\} \tag{2.10}$$

is bounded.

Let T be a Ritt operator and let $\gamma \in (0, \frac{\pi}{2})$ be such that the spectral inclusion (2.7) holds true and the set (2.8) is bounded. Then $A = I - T$ is a sectorial operator. Indeed $1 - B_\gamma \subset \Sigma_\gamma$ and $zR(z, A) = \big((1-z)-1\big)R(1-z, T)$ for any $z \notin \overline{\Sigma_\gamma}$. Hence for $\theta = \gamma$, the set (2.10) is bounded. Thus for any $\alpha > 0$, one can consider the fractional power $(I-T)^\alpha$. We refer e.g. to [20, Chap. 3] for various definitions of these (bounded) operators and their basic properties. Fractional powers of Ritt operators can be expressed by a natural Dunford-Riesz functional calculus formula. Indeed it was observed in [30] that for any polynomial φ, we have

$$\varphi(T)(I-T)^\alpha \,=\, \frac{1}{2\pi i}\int_{\partial B_\gamma} \varphi(\lambda)(1-\lambda)^\alpha R(\lambda, T)\, d\lambda\,, \tag{2.11}$$

where the countour ∂B_γ is oriented counterclockwise.

P. Vitse proved in [52] that if $T\colon X \to X$ is a Ritt operator, then for any integer $N \geq 0$, the set $\{n^N T^{n-1}(I-T)^N \,:\, n \geq 1\}$ is bounded. Our next statement is a continuation of these results.

Proposition 2.8. *Let X be a Banach space and let $T\colon X \to X$ be a Ritt operator (resp. an R-Ritt operator). For any $\alpha > 0$, the set*

$$\big\{n^\alpha (rT)^{n-1}(I-rT)^\alpha \,:\, n \geq 1,\ r \in (0,1]\big\}$$

is bounded (resp. R-bounded).

Proof. We will prove this result in the 'R-Ritt case' only. The 'Ritt case' is similar and simpler. Assume that T is R-Ritt. Applying Lemma 2.7, we let $\gamma \in (0, \frac{\pi}{2})$ be such that (2.7) holds true and the set (2.8) is R-bounded. Let $r \in (0,1]$ and let $\lambda \in \mathbb{C} \setminus B_\gamma$, with $\lambda \neq 1$. Then $\frac{\lambda}{r} \in \mathbb{C} \setminus B_\gamma$ hence $\frac{\lambda}{r}$ belongs to the resolvent set of T and we have

$$(\lambda - 1)R(\lambda, rT) = \frac{\lambda - 1}{\lambda - r}\Big(\frac{\lambda}{r} - 1\Big)R\Big(\frac{\lambda}{r}, T\Big).$$

Since the set

$$\Big\{\frac{\lambda - 1}{\lambda - r} \,:\, \lambda \in \mathbb{C} \setminus B_\gamma,\ \lambda \neq 1,\ r \in (0,1]\Big\}$$

is bounded, it follows from the above formula that the set

$$\big\{(\lambda - 1)R(\lambda, rT) \,:\, \lambda \in \mathbb{C} \setminus B_\gamma,\ \lambda \neq 1,\ r \in (0,1]\big\} \tag{2.12}$$

is R-bounded.

The boundary ∂B_γ is the juxtaposition of the segment Γ_+ going from 1 to $1 - \cos(\gamma)e^{-i\gamma}$, of the segment Γ_- going from $1 - \cos(\gamma)e^{i\gamma}$ to 1 and of the curve Γ_0 going from $1 - \cos(\gamma)e^{-i\gamma}$ to $1 - \cos(\gamma)e^{i\gamma}$ counterclockwise along the circle of center 0 and radius $\sin\gamma$.

Consider a fixed number $\alpha > 0$. For any integer $n \geq 1$ and any $r \in (0,1]$, we have

$$(rT)^{n-1}(I-rT)^\alpha \,=\, \frac{1}{2\pi i}\int_{\partial B_\gamma} \lambda^{n-1}(1-\lambda)^\alpha R(\lambda, rT)\, d\lambda$$

by applying (2.11) to rT. Hence we may write

$$n^\alpha (rT)^{n-1}(I-rT)^\alpha \,=\, \frac{-n^\alpha}{2\pi i}\int_{\partial B_\gamma} \lambda^{n-1}(1-\lambda)^{\alpha-1}(\lambda-1)R(\lambda,rT)\,d\lambda\,.$$

According to the R-boundedness of the set (2.12) and Lemma 2.1, it therefore suffices to show that the integrals

$$I_n \,=\, n^\alpha \int_{\partial B_\gamma} |\lambda|^n |1-\lambda|^{\alpha-1}\,|d\lambda|$$

are uniformly bounded (for n varying in $\mathbb{N}$). Let us decompose each of these integrals as $I_n = I_{n,0} + I_{n,+} + I_{n,-}$, with

$$I_{n,0} = n^\alpha \int_{\Gamma_0} \cdots |d\lambda|\,, \quad I_{n,+} = n^\alpha \int_{\Gamma_+} \cdots |d\lambda|\,, \quad \text{and} \quad I_{n,-} = n^\alpha \int_{\Gamma_-} \cdots |d\lambda|\,.$$

For $\lambda \in \Gamma_0$, we both have

$$\cos\gamma \le |1-\lambda| \le 2 \qquad \text{and} \qquad |\lambda| = \sin\gamma.$$

Since the sequence $\big(n^\alpha(\sin\gamma)^n\big)_{n\ge1}$ is bounded, this readily implies that the sequence $(I_{n,0})_{n\ge1}$ is bounded.

Let us now estimate $I_{n,+}$. For any $t \in [0,\cos\gamma]$, we have $t^2 \le t\cos\gamma$ hence

$$\big|1-te^{-i\gamma}\big|^2 \,=\, 1+t^2-2t\cos\gamma \le 1-t\cos\gamma.$$

Hence

$$I_{n,+} = n^\alpha \int_0^{\cos\gamma} \big|1-te^{-i\gamma}\big|^n t^{\alpha-1}\,dt \;\le n^\alpha \int_0^{\cos\gamma} \big(1-t\cos\gamma\big)^{\frac{n}{2}} t^{\alpha-1}\,dt\,.$$

Changing t into $s = t\cos\gamma$ and using the inequality $1-s \le e^{-s}$, we deduce that

$$I_{n,+} \le \,\frac{n^\alpha}{(\cos\gamma)^\alpha}\int_0^{\cos^2\gamma} s^{\alpha-1}\,e^{-\frac{sn}{2}}\,ds\,.$$

This yields (changing s into $u = \frac{sn}{2}$)

$$I_{n,+} \le \,\frac{2^\alpha}{(\cos\gamma)^\alpha}\int_0^{\infty} u^{\alpha-1}e^{-u}\,du\,.$$

Thus the sequence $(I_{n,+})_{n\ge1}$ is bounded. Since $I_{n,-} = I_{n,+}$, this completes the proof of the boundedness of $(I_n)_{n\ge1}$. □

3. Equivalence of square functions.

Throughout the next two sections, we fix a measure space (Ω,μ) and a number $1<p<\infty$. We shall deal with operators acting on the Banach space $X = L^p(\Omega)$. We start with a precise definition of (1.1) and (1.3) and a few comments.

Let $T\colon L^p(\Omega) \to L^p(\Omega)$ be a bounded operator and let $x \in L^p(\Omega)$. Let us consider

$$x_k = k^{\frac12}\big(T^k(x) - T^{k-1}(x)\big)$$

for any $k \ge 1$. If the sequence $(x_k)_{k\ge1}$ belongs to the space $L^p(\Omega;\ell^2)$, then $\|x\|_{T,1}$ is defined as the norm of $(x_k)_{k\ge1}$ in that space. Otherwise, we set $\|x\|_{T,1} = \infty$. If T is a Ritt operator,

then the quantities $\|x\|_{T,\alpha}$ are defined in a similar manner for any $\alpha > 0$. In particular, $\|x\|_{T,\alpha}$ can be infinite.

These square functions are natural discrete analogs of the square functions asociated to sectorial operators (see [12] and the survey paper [31]).

Assume that T is a Ritt operator. Then T is power bounded hence by the Mean Ergodic Theorem (see e.g. [25, Subsection 2.1.1]), we have a direct sum decomposition

$$L^p(\Omega) \,=\, \mathrm{Ker}(I-T) \oplus \overline{\mathrm{Ran}(I-T)}, \tag{3.1}$$

where $\mathrm{Ker}(\cdot\,)$ and $\mathrm{Ran}(\cdot\,)$ denote the kernel and the range, respectively. For any $\alpha > 0$, we have $\mathrm{Ker}\big((I-T)^\alpha\big) = \mathrm{Ker}(I-T)$. This implies that

$$\|x\|_{T,\alpha} = 0 \iff x \in \mathrm{Ker}(I-T). \tag{3.2}$$

Given any $\alpha > 0$, a general question is to determine whether $\|x\|_{T,\alpha} < \infty$ for any x in $L^p(\Omega)$. It is easy to check, using the Closed graph Theorem, that this finiteness property is equivalent to the existence of a constant $C \geq 0$ such that

$$\|x\|_{T,\alpha} \leq C\|x\|_{L^p}, \qquad x \in L^p(\Omega). \tag{3.3}$$

In [30], the second named author established the following connection between the boundedness of discrete square functions and functional calculus properties.

Theorem 3.1. *([30]) Let $T\colon L^p(\Omega) \to L^p(\Omega)$ be a Ritt operator, with $1 < p < \infty$. The following assertions are equivalent.*

(i) *The operator T and its adjoint $T^*\colon L^{p'}(\Omega) \to L^{p'}(\Omega)$ both satisfy uniform estimates*
$$\|x\|_{T,1} \lesssim \|x\|_{L^p} \qquad \text{and} \qquad \|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}}$$
for $x \in L^p(\Omega)$ and $y \in L^{p'}(\Omega)$.

(ii) *There exists an angle $0 < \gamma < \frac{\pi}{2}$ and a constant $K \geq 0$ such that*
$$\|\varphi(T)\| \,\leq\, K\,\|\varphi\|_{H^\infty(B_\gamma)}$$
for any $\varphi \in \mathcal{P}$.

(iii) *The operator T is R-Ritt and there exists an angle $0 < \theta < \pi$ such that $I - T$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus.*

Besides [30], we refer to [12, 26, 32, 37] for general information on $H^\infty(\Sigma_\theta)$ functional calculus for sectorial operators.

The main purpose of this section is to show that if T is R-Ritt, then the square functions $\|\ \|_{T,\alpha}$ are pairwise equivalent. Thus the existence of an estimate (3.3) does not depend on $\alpha > 0$. This result (Theorem 3.3 below) is a discrete analog of the equivalence of square functions associated to R-sectorial operators, as established in [33].

We start with preliminary results which allow to estimate square functions $\|x\|_{T,\alpha}$ by means of approximation processes.

Lemma 3.2. *Assume that $T\colon L^p(\Omega) \to L^p(\Omega)$ is a Ritt operator, and let $\alpha > 0$.*

(1) *For any operator $V\colon L^p(\Omega) \to L^p(\Omega)$ such that $VT = TV$ and any $x \in L^p(\Omega)$, we have*
$$\|V(x)\|_{T,\alpha} \leq \|V\|\|x\|_{T,\alpha}.$$

(2) *For any* $x \in \mathrm{Ran}(I-T)$, *we have* $\|x\|_{T,\alpha} < \infty$.
(3) *Let* $\nu \geq \alpha+1$ *be an integer and let* $x \in \mathrm{Ran}\big((I-T)^\nu\big)$. *Then*

$$\|x\|_{T,\alpha} \,=\, \lim_{r\to 1^-} \|x\|_{rT,\alpha}.$$

Proof. (1): Consider $V \in B\big(L^p(\Omega)\big)$. As is well-known, the tensor product $V \otimes I_{\ell^2}$ extends to a bounded operator $V\overline{\otimes}I_{\ell^2}\colon L^p(\Omega;\ell^2) \longrightarrow L^p(\Omega;\ell^2)$, with $\|V\overline{\otimes}I_{\ell^2}\| = \|V\|$. Assume that $VT = TV$ and let x be such that $\|x\|_{T,\alpha} < \infty$. Then we have

$$\big(k^{\alpha-\frac{1}{2}}T^{k-1}(I-T)^\alpha\big(V(x)\big)\big)_{k\geq 1} \,=\, V\overline{\otimes}I_{\ell^2}\Big[\big(k^{\alpha-\frac{1}{2}}T^{k-1}(I-T)^\alpha(x)\big)_{k\geq 1}\Big],$$

and the result follows at once.

(2): Assume that $x = (I-T)x'$ for some $x' \in L^p(\Omega)$. By Proposition 2.8, there exists a constant C such that

$$\begin{aligned}\sum_{k=1}^\infty \big\|k^{\alpha-\frac{1}{2}}T^{k-1}(I-T)^\alpha(x)\big\|_{L^p} \,&=\, \sum_{k=1}^\infty k^{\alpha-\frac{1}{2}}\big\|T^{k-1}(I-T)^{\alpha+1}(x')\big\|_{L^p}\\ &\leq\, \|x'\|_{L^p}\sum_{k=1}^\infty k^{\alpha-\frac{1}{2}}\,\frac{C}{k^{\alpha+1}}\\ &\leq\, C\,\|x'\|_{L^p}\sum_{k=1}^\infty k^{-\frac{3}{2}} \,<\infty.\end{aligned}$$

This implies that $\big(k^{\alpha-\frac{1}{2}}T^{k-1}(I-T)^\alpha(x)\big)_{k\geq 1}$ belongs to $L^p(\Omega;\ell^2)$.

(3): It is clear that $(I-rT)^\alpha \to (I-T)^\alpha$ when $r\to 1^-$. Assume that $x \in \mathrm{Ran}\big((I-T)^\nu\big)$. Arguing as in part (2) we find that the sequence $\big(k^{\alpha-\frac{1}{2}}T^{k-1}(x)\big)_{k\geq 1}$ belongs to $L^p(\Omega;\ell^2)$. Then arguing as in part (1), we obtain that

$$\Big\|\Big(k^{\alpha-\frac{1}{2}}T^{k-1}\big((I-rT)^\alpha-(I-T)^\alpha\big)(x)\Big)_{k\geq 1}\Big\|_{L^p(\ell^2)} \longrightarrow\, 0$$

when $r\to 1^-$. This implies the convergence result. □

Theorem 3.3. *Assume that* $T\colon L^p(\Omega)\to L^p(\Omega)$ *is an R-Ritt operator. Then for any* $\alpha,\beta > 0$, *we have an equivalence*

$$\|x\|_{T,\alpha} \,\approx\, \|x\|_{T,\beta}, \qquad x \in L^p(\Omega).$$

Proof. We fix $\gamma > 0$ such that $\alpha+\gamma$ is an integer $N \geq 1$. For any integer $k \geq 1$, we define the complex number

$$c_k = \frac{k(k+1)\cdots(k+N-2)}{k^{\alpha-\frac{1}{2}}},$$

with the convention that $c_k = \frac{1}{k^{\alpha-\frac{1}{2}}}$ if $N=1$. For any $z\in\mathbb{D}$, we have

$$\sum_{k=1}^\infty k(k+1)\cdots(k+N-2)z^{k-1} \,=\, \frac{(N-1)!}{(1-z)^N}.$$

Hence

$$\sum_{k=1}^{\infty} c_k k^{\alpha-\frac{1}{2}} z^{2k-2}(1-z^2)^N \;=\; \sum_{k=1}^{\infty} k(k+1)\cdots(k+N-2)(z^2)^{k-1}(1-z^2)^N \;=\; (N-1)!.$$

Since the operator T is power bounded, we deduce that for every $r \in (0,1)$ we have

$$\sum_{k=1}^{\infty} c_k k^{\alpha-\frac{1}{2}} (rT)^{2k-2}\big(I-(rT)^2\big)^N \;=\; (N-1)!I,$$

the series being absolutely convergent. Since $(I+rT)^N$ is invertible, this yields

$$\sum_{k=1}^{\infty} c_k (rT)^{k-1}(I-rT)^{\gamma} k^{\alpha-\frac{1}{2}} (rT)^{k-1}(I-rT)^{\alpha} = (N-1)!(I+rT)^{-N}.$$

Let $x \in L^p(\Omega)$. For any integer $m \geq 1$ and any $r \in (0,1)$, we let

$$y_m(r) \;=\; (N-1)!(I+rT)^{-N} m^{\beta-\frac{1}{2}} (rT)^{m-1}(I-rT)^{\beta} x.$$

Then it follows from the above identity that

$$y_m(r) \;=\; \sum_{k=1}^{\infty} c_k m^{\beta-\frac{1}{2}} (rT)^{m+k-2}(I-rT)^{\beta+\gamma} \cdot k^{\alpha-\frac{1}{2}} (rT)^{k-1}(I-rT)^{\alpha} x.$$

For any $n \geq 1$, we consider the partial sum

$$y_{m,n}(r) \;=\; \sum_{k=1}^{n} c_k m^{\beta-\frac{1}{2}} (rT)^{m+k-2}(I-rT)^{\beta+\gamma} \cdot k^{\alpha-\frac{1}{2}} (rT)^{k-1}(I-rT)^{\alpha} x,$$

and we have $y_{m,n}(r) \to y_m(r)$ when $n \to \infty$.

Let us write

(3.4)

$$c_k m^{\beta-\frac{1}{2}} (rT)^{m+k-2}(I-rT)^{\beta+\gamma} \;=\; \frac{m^{\beta-\frac{1}{2}} c_k}{(m+k-1)^{\gamma+\beta}} \big[(m+k-1)^{\gamma+\beta}(rT)^{m+k-2}(I-rT)^{\beta+\gamma}\big]$$

for any $m,k \geq 1$. Since $c_k \sim_{+\infty} k^{\gamma-\frac{1}{2}}$, there exists a positive constant K such that

$$\frac{m^{\beta-\frac{1}{2}} c_k}{(m+k-1)^{\gamma+\beta}} \leq K\, \frac{m^{\beta-\frac{1}{2}} k^{\gamma-\frac{1}{2}}}{(m+k)^{\gamma+\beta}}$$

for any $m,k \geq 1$. It therefore follows from Proposition 2.3 and Lemma 2.4 that the matrix

$$\left[\frac{m^{\beta-\frac{1}{2}} c_k}{(m+k-1)^{\gamma+\beta}}\right]_{m,k\geq 1}$$

represents an element of $B(\ell^2)$. Moreover, by Proposition 2.8, the set

$$F \;=\; \big\{(m+k-1)^{\gamma+\beta}(rT)^{m+k-2}(I-rT)^{\gamma+\beta} \;:\; m,k \geq 1,\; r \in (0,1]\big\}$$

is R-bounded. Hence by (2.1), (3.4) and Proposition 2.6, we get to an estimate

$$\left\| \left(\sum_{m=1}^{M} \big|y_{m,n}(r)\big|^2 \right)^{\frac{1}{2}} \right\|_{L^p} \lesssim \left\| \left(\sum_{k=1}^{\infty} k^{2\alpha-1} \big|(rT)^{k-1}(I-rT)^{\alpha} x\big|^2 \right)^{\frac{1}{2}} \right\|_{L^p}$$

for any integer $M \geq 1$. Passing to the limit, we deduce that

$$\Big\| \Big(\sum_{m=1}^{\infty} \big| y_m(r) \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p} \lesssim \Big\| \Big(\sum_{k=1}^{\infty} k^{2\alpha-1} \big| (rT)^{k-1} (I - rT)^{\alpha} x \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p}.$$

Since the set $\big\{ (N-1)!^{-1} (I + rT)^N \, : \, r \in (0,1) \big\}$ is bounded, we finally obtain that

$$\|x\|_{rT,\beta} \lesssim \|x\|_{rT,\alpha}.$$

It is crucial to note that in this estimate, the majorizing constant hidden in the symbol $\lesssim$ does not depend on $r \in (0,1)$.

Now let ν be an integer such that $\nu \geq \alpha + 1$ and $\nu \geq \beta + 1$. Applying Lemma 3.2 (3), we deduce a uniform estimate

$$\|x\|_{T,\beta} \lesssim \|x\|_{T,\alpha}$$

for $x \in \mathrm{Ran}\big((I-T)^{\nu} \big)$. Next for any integer $m \geq 0$, set

$$\Lambda_m = \frac{1}{m+1} \sum_{k=0}^{m} (I - T^k).$$

It is clear that Λ_m^{ν} maps X into $\mathrm{Ran}\big((I-T)^{\nu} \big)$. Hence we actually have a uniform estimate

$$\big\| \Lambda_m^{\nu}(x) \big\|_{T,\beta} \lesssim \big\| \Lambda_m^{\nu}(x) \big\|_{T,\alpha}, \qquad x \in X, \ m \geq 1.$$

Since T is power bounded, the sequence $(\Lambda_m)_{m \geq 0}$ is bounded. Applying Lemma 3.2 (1), we deduce a further uniform estimate

$$\big\| \Lambda_m^{\nu}(x) \big\|_{T,\beta} \lesssim \|x\|_{T,\alpha}, \qquad x \in X, \ m \geq 1.$$

Equivalently, we have

$$\Big\| \Big(\sum_{k=1}^{l} k^{2\beta-1} \big| T^{k-1} (I-T)^{\beta} \Lambda_m^{\nu}(x) \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p} \lesssim \|x\|_{T,\alpha}, \qquad x \in X, \ m \geq 1, \ l \geq 1.$$

For any $x \in \overline{\mathrm{Ran}(I-T)}$, $\Lambda_m(x) \to x$ and hence $\Lambda_m^{\nu}(x) \to x$ when $m \to \infty$. Hence passing to the limit in the above inequaliy, we obtain a uniform estimate $\|x\|_{T,\beta} \lesssim \|x\|_{T,\alpha}$ for x in $\overline{\mathrm{Ran}(I-T)}$. Switching the roles of α and β, this shows that $\| \ \|_{T,\beta}$ and $\| \ \|_{T,\alpha}$ are equivalent on the space $\overline{\mathrm{Ran}(I-T)}$. Moreover $\| \ \|_{T,\beta}$ and $\| \ \|_{T,\alpha}$ vanish on $\mathrm{Ker}(I-T)$ by (3.2). Appealing to the direct sum decomposition (3.1), we finally obtain that $\| \ \|_{T,\beta}$ and $\| \ \|_{T,\alpha}$ are equivalent on $L^p(\Omega)$. □

The techniques developed so far in this paper allow us to prove the following proposition, which complements Theorem 3.1. For a Ritt operator T, we let P_T denote the projection onto $\mathrm{Ker}(I-T)$ which vanishes on $\overline{\mathrm{Ran}(I-T)}$ (recall (3.1)).

Proposition 3.4. *Let $T \colon L^p(\Omega) \to L^p(\Omega)$ be a Ritt operator, with $1 < p < \infty$. Then the condition (i) in Theorem 3.1 is equivalent to:*

(i)' *We have an equivalence*

$$\|x\|_{L^p} \approx \|P_T(x)\|_{L^p} + \|x\|_{T,1}$$

for $x \in L^p(\Omega)$.

Proof. That (i) implies (i)' was proved in [34, Rem. 3.4] in the case when T is 'contractively regular'. The proof in our present case is the same.

Assume (i)'. Let $y \in L^{p'}(\Omega)$. We consider a finite sequence $(x_k)_{k\geq 1}$ in $L^p(\Omega)$ and we set

$$x \,=\, \sum_k k^{\frac{1}{2}} T^{k-1}(I-T)x_k\,.$$

Then

$$\left|\sum_k \big\langle k^{\frac{1}{2}}(T^*)^{k-1}(I-T^*)y, x_k\big\rangle\right| \,=\, |\langle y, x\rangle| \,\leq\, \|x\|_{L^p}\|y\|_{L^{p'}}.$$

Moreover $x \in \mathrm{Ran}(I-T)$ hence applying (i)', we deduce

$$\left|\sum_k \big\langle k^{\frac{1}{2}}(T^*)^{k-1}(I-T^*)y, x_k\big\rangle\right| \,\lesssim\, \|y\|_{L^{p'}}\|x\|_{T,1}.$$

We will now show an estimate

$$\|x\|_{T,1} \,\lesssim\, \left\|\left(\sum_k |x_k|^2\right)^{\frac{1}{2}}\right\|_{L^p}. \tag{3.5}$$

Then passing to the supremum over all finite sequences $(x_k)_{k\geq 1}$ in the unit ball of $L^p(\Omega;\ell^2)$, we deduce that $\|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}}$.

To show (3.5), first note that for any integer $m \geq 1$, we may write

$$m^{\frac{1}{2}} T^{m-1}(I-T)x \,=\, \sum_k \frac{m^{\frac{1}{2}} k^{\frac{1}{2}}}{(m+k)^2}\,(m+k)^2 T^{m+k-2}(I-T)^2 x_k\,.$$

Second according to [30], the assumption (i)' implies that T is an R-Ritt operator. Hence by Proposition 2.8, the set $\{(m+k)^2 T^{m+k-2}(I-T)^2 \,:\, m,k \geq 1\}$ is R-bounded. Therefore applying Propositions 2.3 and 2.6 we obtain (3.5). □

4. Loose dilations

We will focus on the following notion of dilation for L^p-operators.

Definition 4.1. *Let $T\colon L^p(\Omega) \to L^p(\Omega)$ be a bounded operator. We say that it admits a loose dilation if there exist a measure space $(\widetilde{\Omega}, \widetilde{\mu})$, two bounded maps $J\colon L^p(\Omega) \to L^p(\widetilde{\Omega})$ and $Q\colon L^p(\widetilde{\Omega}) \to L^p(\Omega)$, as well as an isomorphism $U\colon L^p(\widetilde{\Omega}) \to L^p(\widetilde{\Omega})$ such that $\big\{U^n \,:\, n \in \mathbb{Z}\big\}$ is bounded and*

$$T^n = QU^nJ, \qquad n \geq 0.$$

That notion is strictly weaker than the following more classical one.

Remark 4.2. We say that a bounded operator $T\colon L^p(\Omega) \to L^p(\Omega)$ admits a *strict dilation* if there exist a measure space $(\widetilde{\Omega}, \widetilde{\mu})$, two contractions $J\colon L^p(\Omega) \to L^p(\widetilde{\Omega})$ and $Q\colon L^p(\widetilde{\Omega}) \to L^p(\Omega)$, as well as an isometric isomorphism $U\colon L^p(\widetilde{\Omega}) \to L^p(\widetilde{\Omega})$ such that $T^n = QU^nJ$ for any $n \geq 0$.

This strict dilation property implies that T is a contraction and that J and Q^* are both isometries.

Conversely in the case $p = 2$, Nagy's dilation Theorem (see e.g. [51, Chapter 1]) ensures that any contraction $L^2(\Omega) \to L^2(\Omega)$ admits a strict dilation.

Next, assume that $1 < p \neq 2 < \infty$. Then it follows from [1, 2, 10, 42] that $T\colon L^p(\Omega) \to L^p(\Omega)$ admits a strict dilation if and only if there exists a positive contraction $S\colon L^p(\Omega) \to L^p(\Omega)$ such that $|T(x)| \leq S(|x|)$ for any $x \in L^p(\Omega)$.

Except for $p = 2$ (see Remark 4.3 below), there is no similar description of operators admitting a loose dilation. The general issue behind our investigation is to try to characterize the L^p-operators which satisfy this property. Theorem 4.8 below gives a satisfactory answer for the class of Ritt operators.

Remark 4.3. Let H be a Hilbert space, let $T\colon H \to H$ be a bounded operator and let us say that T admits a loose dilation if there exist a Hilbert space K, two bounded maps $J\colon H \to K$ and $Q\colon K \to H$, and an isomorphism $U\colon K \to K$ such that $\{U^n : n \in \mathbb{Z}\}$ is bounded and $T^n = QU^nJ$ for any $n \geq 0$. Then this property is equivalent to T being similar to a contraction.

Indeed assume that there exists an isomorphism $V \in B(H)$ such that $V^{-1}TV$ is a contraction. By Nagy's dilation Theorem, that contraction admits a unitary dilation. In other words, there is a unitary U on a Hilbert space K containing H, such that $(V^{-1}TV)^n = qU^nj$ for any $n \geq 0$, where $j\colon H \to K$ is the canonical inclusion and $q = j^*$ is the corresponding orthogonal projection. We obtain the loose dilation property of T by taking $J = jV^{-1}$ and $Q = Vq$.

The converse uses the notion of complete polynomial boundedness, for which we refer to [40, 41]. Assume that T admits a loose dilation. Using [41, Cor. 9.4] and elementary arguments, we obtain that T is completely polynomially bounded. Hence it is similar to a contraction by [40, Cor. 3.5].

According to the above result, the rest of this section is significant only in the case $1 < p \neq 2 < \infty$.

Let $S\colon \ell^p_{\mathbb{Z}} \to \ell^p_{\mathbb{Z}}$ denote the natural shift operator given by $S\big((t_k)_{k\in\mathbb{Z}}\big) = \big((t_{k-1})_{k\in\mathbb{Z}}\big)$. For any φ in $\mathcal{P}$ (the algebra of complex polynomials), we set

$$\|\varphi\|_p = \big\|\varphi(S)\big\|_{B(\ell^p_{\mathbb{Z}})}. \tag{4.1}$$

We recall that if φ is given by $\varphi(z) = \sum_{k\geq 0} d_k z^k$, then $\varphi(S)$ is the convolution operator (with respect to the group $\mathbb{Z}$) associated to the sequence $(d_k)_{k\in\mathbb{Z}}$. Alternatively, $\varphi(S)$ is the Fourier multiplier associated to the restriction of φ to the unitary group $\mathbb{T}$. We refer the reader to [17] for some elementary background on Fourier multiplier theory.

Let us decompose $(0, \pi)$ dyadically into the following family $(I_j)_{j\in\mathbb{Z}}$ of intervals:

$$I_j = \begin{cases} \left[\pi - \frac{\pi}{2^{j+1}}, \pi - \frac{\pi}{2^{j+2}}\right) & \text{if } j \geq 0 \\ \left[2^{j-1}\pi, 2^j\pi\right) & \text{if } j < 0. \end{cases}$$

Then we denote by Δ_j the corresponding arcs of $\mathbb{T}$:

$$\Delta_j = \{e^{it} : t \in -I_j \cup I_j\}.$$

We will use the following version of the Marcinkiewicz multiplier theorem (see [6, Thm. 4.3] and also [17]).

Theorem 4.4. *Let $1 < p < \infty$. Let $\phi \in L^\infty(\mathbb{T})$ and assume that ϕ has uniformly bounded variations over the $(\Delta_j)_{j\in\mathbb{Z}}$. Then ϕ induces a bounded Fourier multiplier $M_\phi \colon \ell^p_{\mathbb{Z}} \to \ell^p_{\mathbb{Z}}$ and we have*

$$\big\|M_\phi\big\|_{B(\ell^p_{\mathbb{Z}})} \le C_p \Big(\|\phi\|_{L^\infty(\mathbb{T})} + \sup\big\{\mathrm{var}\,(\phi, \Delta_j) \,:\, j \in \mathbb{Z}\big\}\Big),$$

where $\mathrm{var}\,(\phi, \Delta_j)$ is the usual variation of ϕ over Δ_j and the constant C_p only depends on p.

For convenience, Definition 4.5 and Proposition 4.7 below are given for an arbitrary Banach space X, although we are mostly interested in the case when X is an L^p-space.

Definition 4.5. *We say that a bounded operator $T \colon X \to X$ is p-polynomially bounded is there exists a constant $C \ge 1$ such that*

$$\big\|\varphi(T)\big\| \le C\|\varphi\|_p \tag{4.2}$$

for any complex polynomial φ.

The following connection with dilations is well-known.

Proposition 4.6. *If $T \colon L^p(\Omega) \to L^p(\Omega)$ admits a loose dilation, then it is p-polynomially bounded.*

Proof. Assume that T satisfies the dilation property given by Definition 4.1. Then for any $\varphi \in \mathcal{P}$, we have $\varphi(T) = Q\varphi(U)J$, hence

$$\big\|\varphi(T)\big\| \le \|Q\|\|J\|\big\|\varphi(U)\big\|.$$

Moreover by the transference principle (see [11, Thm. 2.4]), $\big\|\varphi(U)\big\| \le K^2\|\varphi\|_p$, where $K \ge 1$ is any constant such that $\|U^n\| \le K$ for any integer n. This yields the result. □

We will see in Section 5 that the converse of that proposition does not hold true.

The above proof shows that if $T \colon L^p(\Omega) \to L^p(\Omega)$ admits a strict dilation, then $\|\varphi(T)\| \le \|\varphi\|_p$ for any $\varphi \in \mathcal{P}$, a very classical fact. The famous Matsaev Conjecture asks whether this inequality holds for any L^p-contraction T (even those with no strict dilation). This was disproved very recently by Drury in the case $p = 4$ [16]. It is unclear whether there exists an L^p-contraction T satisfying $\|\varphi(T)\| \le \|\varphi\|_p$ for any $\varphi \in \mathcal{P}$, without admitting a strict dilation.

Proposition 4.7. *Let $T \colon X \to X$ be a p-polynomially bounded operator. Then $I - T$ is sectorial and for any $\theta \in \big(\frac{\pi}{2}, \pi\big)$, it admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus.*

Proof. Since T is p-polynomially bounded, it is power bounded hence $\sigma(T) \subset \overline{\mathbb{D}}$. We can thus define $\varphi(T)$ for any rational function with poles outside $\overline{\mathbb{D}}$. Furthermore (4.2) holds as well for such functions, by approximation.

We fix two numbers $\frac{\pi}{2} < \theta < \theta' < \pi$ and we let (see Figure 2):

$$\mathbb{D}_\theta = D\big(-i\cot(\theta), \tfrac{1}{\sin(\theta)}\big) \cup D\big(i\cot(\theta), \tfrac{1}{\sin(\theta)}\big).$$



Clearly $\mathbb{D}_\theta$ contains $\mathbb{D}$. For any $t \in (-\pi, 0) \cup (0, \pi)$, let $r(t)$ denote the radius of the largest open disc centered at e^{it} and included in $\mathbb{D}_\theta$. If t is positive and small enough, we have

$$\begin{aligned} r(t) &= \frac{1}{\sin(\theta)} - \left|e^{it} + i\cot(\theta)\right| \\ &= \frac{1}{\sin(\theta)} - \sqrt{\cos^2(t) + \big(\sin(t) + \cot(\theta)\big)^2} \\ &= \frac{1}{\sin(\theta)}\Big(1 - \sqrt{1 + 2\sin(t)\sin(\theta)\cos(\theta)}\Big) \\ &= -\cos(\theta)t + \frac{1}{2}\big(\sin(\theta)\cos^2(\theta)\big)t^2 + O(t^3). \end{aligned}$$

Consequently, we have $r(t) > -\cos(\theta)t$ for $t > 0$ small enough. We deduce that if $j < 0$ with $|j|$ large enough and $t \in I_j$, we have

$$D\Big(e^{it}, -\cos(\theta)\frac{\pi}{2^{|j|+1}}\Big) \subset \mathbb{D}_\theta. \tag{4.3}$$

The same holds for $t \in -I_j$. Moreover the intervals I_j and I_{-j} of the dyadic decomposition have length equal to $\frac{\pi}{2^{|j|+1}}$. Hence for any rational function φ with poles outside $\overline{\mathbb{D}_\theta}$ and any $j < 0$ with $|j|$ large enough, we obtain that

$$\begin{aligned} \operatorname{var}\big(\varphi_{|\mathbb{T}}, \Delta_j\big) &= \int_{-I_j \cup I_j} \big|\varphi'(e^{it})\big|\, dt \\ &\le \int_{-I_j \cup I_j} \frac{\|\varphi\|_{H^\infty(\mathbb{D}_\theta)}}{-\cos(\theta)\frac{\pi}{2^{|j|+1}}}\, dt \qquad \text{by (4.3) and Cauchy's inequalities,} \\ &\le \frac{\pi}{2^{|j|}} \cdot \frac{\|\varphi\|_{H^\infty(\mathbb{D}_\theta)}}{-\cos(\theta)\frac{\pi}{2^{|j|+1}}} = \frac{2\pi\|\varphi\|_{H^\infty(\mathbb{D}_\theta)}}{-\cos(\theta)}. \end{aligned}$$

We have a similar result if $j \geq 0$ and large enough. Applying Theorem 4.4, we deduce a uniform estimate

$$\big\|\varphi(S)\big\|_{B(\ell^p_{\mathbb{Z}})} \lesssim \|\varphi\|_{H^\infty(\mathbb{D}_\theta)}.$$

Combining with (4.2) -as explained at the beginning of this proof- we obtain the existence of a constant $K \geq 0$ such that for any rational function φ with poles outside $\overline{\mathbb{D}_\theta}$,

$$\big\|\varphi(T)\big\|_{B(X)} \leq K\|\varphi\|_{H^\infty(\mathbb{D}_\theta)}.$$

Note that we have the following inclusion:

$$1 - \mathbb{D}_\theta \subset \Sigma_\theta.$$

Then let $\mathcal{R}_{\theta'}$ be the algebra of all rational functions with poles outside $\overline{\Sigma_{\theta'}}$ and with a nonpositive degree. We deduce from above that for any $f \in \mathcal{R}_{\theta'}$,

$$\big\|f(I-T)\big\| \leq K\big\|f(1-\cdot\,)\big\|_{H^\infty(\mathbb{D}_\theta)} \leq K\|f\|_{H^\infty(\Sigma_\theta)}.$$

According to [32, Prop. 2.10], this readily implies that $I - T$ is sectorial and admits a bounded $H^\infty(\Sigma_{\theta'})$ functional calculus. □

Theorem 4.8. *Let $T\colon L^p(\Omega) \to L^p(\Omega)$ be a bounded operator, with $1 < p < \infty$. The following assertions are equivalent.*

(i) *The operator T and its adjoint $T^*\colon L^{p'}(\Omega) \to L^{p'}(\Omega)$ both satisfy uniform estimates*

$$\|x\|_{T,1} \lesssim \|x\|_{L^p} \qquad \text{and} \qquad \|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}}$$

for $x \in L^p(\Omega)$ and $y \in L^{p'}(\Omega)$.

(ii) *The operator T is R-Ritt and admits a loose dilation.*

(iii) *The operator T is R-Ritt and p-polynomially bounded.*

Proof. That (ii) implies (iii) follows from Proposition 4.6.

Assume (iii). By Proposition 4.7, $I - T$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus for any $\theta > \frac{\pi}{2}$. Since T is R-Ritt, this implies (i) by Theorem 3.1.

Assume (i). It follows from [30] (see Theorem 3.1) that T is an R-Ritt operator. Thus we only need to establish the dilation property of T. Since T is R-Ritt, Theorem 3.3 ensures that the square functions $\|\ \|_{T,1}$ and $\|\ \|_{T,\frac{1}{2}}$ are equivalent on $L^p(\Omega)$. Likewise, $\|\ \|_{T^*,1}$ and $\|\ \|_{T^*,\frac{1}{2}}$ are equivalent on $L^{p'}(\Omega)$. Consequently the assumption (i) implies the existence of a constant $C \geq 1$ such that

$$\|x\|_{T,\frac{1}{2}} \leq C\|x\|_{L^p} \qquad \text{and} \qquad \|y\|_{T^*,\frac{1}{2}} \leq C\|y\|_{L^{p'}}$$

for any $x \in L^p(\Omega)$ and any $y \in L^{p'}(\Omega)$.

We will use the direct sum decomposition (3.1), as well as the analogous decompostion of $L^{p'}(\Omega)$ corresponding to T^*. According to the above estimates, we may define two bounded maps

$$j_1\colon \overline{\operatorname{Ran}(I-T)} \longrightarrow L^p(\Omega;\ell^2_{\mathbb{Z}}) \qquad \text{and} \qquad j_2\colon \overline{\operatorname{Ran}(I-T^*)} \longrightarrow L^{p'}(\Omega;\ell^2_{\mathbb{Z}})$$

as follows. For any $x \in \overline{\operatorname{Ran}(I-T)}$ and any $y \in \overline{\operatorname{Ran}(I-T^*)}$, we set

$$x_k = T^{k-1}(I-T)^{\frac{1}{2}}x \qquad \text{and} \qquad y_k = (T^*)^{k-1}(I-T^*)^{\frac{1}{2}}y$$

if $k \geq 0$, we set $x_k = 0$ and $y_k = 0$ if $k < 0$. Then we set

$$j_1(x) = (x_k)_{k\in\mathbb{Z}} \qquad \text{and} \qquad j_2(y) = (y_k)_{k\in\mathbb{Z}}.$$

Then we let $J_1 \colon L^p(\Omega) \to L^p(\Omega) \overset{p}{\oplus} L^p(\Omega;\ell^2_{\mathbb{Z}})$ be the linear map taking any $x \in \operatorname{Ker}(I-T)$ to $(x,0)$ and any $x \in \overline{\operatorname{Ran}(I-T)}$ to $(0, j_1(x))$. We define $J_2 \colon L^{p'}(\Omega) \to L^{p'}(\Omega) \overset{p'}{\oplus} L^{p'}(\Omega;\ell^2_{\mathbb{Z}})$ in a similar way.

For any $x \in \overline{\operatorname{Ran}(I-T)}$ and $y \in \overline{\operatorname{Ran}(I-T^*)}$, we have

$$\begin{aligned}\langle J_1x, J_2y\rangle &= \sum_{k=1}^{\infty} \big\langle T^{k-1}(I-T)^{\frac12}x, (T^*)^{k-1}(I-T^*)^{\frac12}y\big\rangle\\ &= \sum_{k=1}^{\infty} \big\langle T^{2(k-1)}(I-T)x, y\big\rangle\\ &= \sum_{k=1}^{\infty} \big\langle T^{2(k-1)}(I-T^2)(I+T)^{-1}x, y\big\rangle.\end{aligned}$$

For any integer $N \geq 1$

$$\sum_{k=1}^{N} T^{2(k-1)}(I-T^2) \;=\; I - T^{2N}.$$

Furthermore, $(I+T)^{-1}x$ belongs to $\overline{\operatorname{Ran}(I-T)}$ and the sequence $(T^n)_{n\geq 0}$ strongly converges to 0 on that subspace of $L^p(\Omega)$. Hence

$$\langle J_1x, J_2y\rangle \;=\; \langle (I+T)^{-1}x, y\rangle.$$

Let $\Theta \colon L^p(\Omega) \to L^p(\Omega)$ be the linear map taking any $x \in \overline{\operatorname{Ran}(I-T)}$ to $(I+T)x$ and any $x \in \operatorname{Ker}(I-T)$ to itself. Then it follows from the above calculation that

$$\Theta J_2^* J_1 \;=\; I_{L^p(\Omega)}. \tag{4.4}$$

Let

$$Z = L^p(\Omega) \overset{p}{\oplus} L^p(\Omega;\ell^2_{\mathbb{Z}}),$$

and let $U \colon Z \to Z$ be the linear map which takes any $x \in L^p(\Omega)$ to itself and any sequence $(x_k)_{k\in\mathbb{Z}}$ in $L^p(\Omega;\ell^2_{\mathbb{Z}})$ to the shifted sequence $(x_{k+1})_{k\in\mathbb{Z}}$. Next let $P \colon Z \to Z$ be the linear map which takes any $x \in L^p(\Omega)$ to itself and any sequence $(x_k)_{k\in\mathbb{Z}}$ in $L^p(\Omega;\ell^2_{\mathbb{Z}})$ to the truncated sequence $(\dots, 0, \dots, 0, x_0, x_1, \dots, x_k, \dots)$. By construction, we have

$$PU^nJ_1 = J_1T^n, \qquad n \geq 0. \tag{4.5}$$

We also have $J_2^*P = J_2^*$ hence setting $J = J_1 \colon L^p(\Omega) \to Z$ and $Q = \Theta J_2^* \colon Z \to L^p(\Omega)$, we deduce from (4.4) and (4.5) that $T^n = QU^nJ$ for any $n \geq 0$. Furthermore, U is an isometric isomorphism on Z. Thus we have established that T satisfies the dilation property stated in Definition 4.1, except that the dilation space is Z instead of being an L^p-space.

It is easy to modify the construction to obtain a dilation through an L^p-space, as follows. First recall that using for example Gaussian variables, one can isometrically represent $\ell^2_{\mathbb{Z}}$ as

a complemented subspace of an L^p-space (see e.g. [43, Chapter 5]). The space Z can be therefore represented as well as a complemented subspace of an L^p-space. Thus we have

$$Z \oplus W \;=\; L^p(\widetilde{\Omega})$$

for an appropriate measure space $(\widetilde{\Omega}, \widetilde{\mu})$ and some Banach space W. Let $J'\colon L^p(\Omega) \to L^p(\widetilde{\Omega})$ be defined by $J'(x) = (J(x), 0)$, let $U'\colon L^p(\widetilde{\Omega}) \to L^p(\widetilde{\Omega})$ be defined by $U'(z,w) = (Uz, w)$ and let $Q'\colon L^p(\widetilde{\Omega}) \to L^p(\Omega)$ be defined by $Q'(z,w) = Q(z)$. Then U' is an isomorphism, $(U'^n)_{n\in\mathbb{Z}}$ is bounded and $Q'U'^nJ' = T^n$ for any $n \geq 0$. □

5. Comparing p-boundedness properties

In this section we will consider an L^p-analog of complete polynomial boundedness going back to [45] (see also [47, Chap. 8]) and give complements to the results obtained in the previous section. In particular we will show the existence of p-polynomially bounded operators $L^p \to L^p$ without any loose dilation.

In the sequel we assume that $1 \leq p < \infty$. Let $n \geq 1$ be an integer. For any vector space V, we let $M_n(V)$ denote the space of $n \times n$ matrices with entries in V. When $V = B(X)$ for some Banach space X, we equip this space with a specific norm, as follows. For any $[T_{ij}]_{1\leq i,j\leq n}$ in $M_n\big(B(X)\big)$, we set

$$\big\|[T_{ij}]\big\|_{p,M_n(B(X))} \;=\; \sup\left\{ \left(\sum_{i=1}^n \Big\| \sum_{j=1}^n T_{ij}(x_j) \Big\|_X^p \right)^{\frac{1}{p}} \;:\; x_1,\dots,x_n \in X,\; \sum_{j=1}^n \|x_j\|_X^p \leq 1 \right\}. \tag{5.1}$$

In other words, we regard $[T_{ij}]$ as an operator $\ell_n^p(X) \to \ell_n^p(X)$ in a natural way and the norm of the matrix is defined as the corresponding operator norm.

Let X, Y be two Banach spaces, let $V \subset B(X)$ be a subspace and let $u\colon V \to B(Y)$ be a linear mapping. We say that u is p-completely bounded if there exists a constant $C \geq 0$ such that

$$\big\|[u(T_{ij})]\big\|_{p,M_n(B(Y))} \;\leq C \big\|[T_{ij}]\big\|_{p,M_n(B(X))}$$

for any $n \geq 1$ and any matrix $[T_{ij}]$ in $M_n(V)$. In this case, we let $\|u\|_{pcb}$ denote the smallest possible C.

Let us regard the vector space $\mathcal{P}$ of all complex polynomials as a subspace of $B(\ell_{\mathbb{Z}}^p)$, by identifying any $\varphi \in \mathcal{P}$ with the operator $\varphi(S)$. Accordingly for any $[\varphi_{ij}]$ in $M_n(\mathcal{P})$, we set

$$\big\|[\varphi_{ij}]\big\|_p \;=\; \big\|[\varphi_{ij}(S)]\big\|_{p,M_n(B(\ell_{\mathbb{Z}}^p))}.$$

This extends (4.1) to matrices. We say that a bounded operator $T\colon Y \to Y$ is p-completely polynomially bounded if the natural mapping $u\colon \mathcal{P} \to B(Y)$ given by $u(\varphi) = \varphi(T)$ is p-completely bounded. This is equivalent to the existence of a constant $C \geq 1$ such that

$$\big\|[\varphi_{ij}(T)]\big\|_{p,M_n(B(Y))} \;\leq C \big\|[\varphi_{ij}]\big\|_p$$

for any matrix $[\varphi_{ij}]$ of complex polynomials.

When $p = 2$ and Y is a Hilbert space, the notions of 2-polynomial boundedness and 2-complete polynomial boundedness correspond to the usual notions of polynomial boundedness and complete polynomial boundedness from [40, 41]. See [41] for the rich connections

with operator space theory. The existence of a polynomially bounded operator on Hilbert space which is not completely polynomially bounded is a major result due to Pisier. Indeed this is the heart of his negative solution to the Halmos problem [46, 47]. We will show that Pisier's construction can be transferred to our L^p-setting.

We start with an elementary result which is obvious when $p = 2$ but requires attention when $p \neq 2$.

Lemma 5.1. *Let $N \geq 1$ be an integer, let H be a Hilbert space and let $\pi\colon B(\ell_N^2) \to B(H)$ be a unital $*$-representation. Then for any $n \geq 1$ and any matrix $[T_{ij}]$ in $M_n\big(B(\ell_N^2)\big)$, we have*

$$\big\|[T_{ij}]\big\|_{p,M_n(B(\ell_N^2))} \leq \big\|[\pi(T_{ij})]\big\|_{p,M_n(B(H))}.$$

Proof. As is well-known, there is a Hilbert space K such that

$$H \simeq \ell_N^2(K), \qquad B(H) \simeq B(\ell_N^2) \otimes B(K),$$

and $\pi(T) = T \otimes I_K$ for any $T \in B(\ell_N^2)$ (see e.g. [13, Cor. III.1.7]). Consider $[T_{ij}]$ in $M_n\big(B(\ell_N^2)\big)$ and $x_1, \ldots, x_n$ in ℓ_N^2. Fix some $e \in K$ with $\|e\| = 1$. Then

$$\begin{aligned}
\sum_i \Big\| \sum_j T_{ij}(x_j) \Big\|_{\ell_N^2}^p &= \sum_i \Big\| \sum_j T_{ij}(x_j) \otimes e \Big\|_{\ell_N^2(K)}^p \\
&= \sum_i \Big\| \sum_j \big[\pi(T_{ij})\big](x_j \otimes e) \Big\|_{\ell_N^2(K)}^p \\
&\leq \big\|[\pi(T_{ij})]\big\|_{p,M_n(B(H))}^p \sum_j \|x_j \otimes e\|_{\ell_N^2(K)}^p \\
&\leq \big\|[\pi(T_{ij})]\big\|_{p,M_n(B(H))}^p \sum_j \|x_j\|_{\ell_N^2}^p ,
\end{aligned}$$

and the result follows at once. □

Proposition 5.2. *Suppose that $1 < p < \infty$. There exists a p-polynomially bounded operator $T\colon L^p\big([0,1]\big) \to L^p\big([0,1]\big)$ which is not p-completely polynomially bounded.*

Proof. We need some background on Pisier's counterexample. We refer to [47, Chap. 9] and [41, Chap. 10] for a detailed exposition of this example and also to the necessary background on Hankel operators on $B(\ell^2(H))$ and their $B(H)$-valued symbols.

We start with a concrete description of a sequence of operators satisfying the so-called canonical anticommutation relations. Let I_2 denote the identity matrix on M_2. For any $k \geq 1$, consider the unital embedding $M_{2^k} \hookrightarrow M_{2^{k+1}} \simeq M_{2^k} \otimes M_2$ given by $A \mapsto A \otimes I_2$. The closure of the union of the resulting increasing sequence $(M_{2^k})_{k\geq 1}$ is a C^*-algebra. Representing it as an algebra of operators, we obtain a Hilbert space H and an embedding

$$\bigcup_{k\geq 1} \uparrow M_{2^k} \subset B(H) \tag{5.2}$$

whose restriction to each M_{2^k} is a unital $*$-representation.

Consider the 2×2 matrices

$$D = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \qquad \text{and} \qquad E = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

For any $k \geq 1$, we set

$$C_k = E^{\otimes(k-1)} \otimes D \ \in M_{2^k},$$

where $E^{\otimes(k-1)}$ denotes the tensor product of E with itself $(k-1)$ times. Then following (5.2) we let $\widetilde{C_k}$ denote this operator regarded as an element of $B(H)$. The distinction between C_k and $\widetilde{C_k}$ may look superfluous. The reason why we need this is that the inclusion providing the identification between C_k and $\widetilde{C_k}$ is a $*$-representation and a priori, $*$-representations are not p-complete isometries (i.e. they do not preserve p-matrix norms). However using Lemma 5.1, we see that for any $m \geq 1$, for any $n \geq 1$ and for any $a_1, \ldots, a_m \in M_n$,

$$\Big\| \sum_{k=1}^m a_k \otimes C_k \otimes I_2^{\otimes(m-k)} \Big\|_{p, M_n(B(\ell_{2^m}^2))} \leq \Big\| \sum_{k=1}^m a_k \otimes \widetilde{C_k} \Big\|_{p, M_n(B(H))}. \tag{5.3}$$

The above sequence of matrices has the following remarkable property (see [47, p. 70]): for any complex numbers $\alpha_1, \ldots, \alpha_m$,

$$\Big\| \sum_{k=1}^m \alpha_k \, C_k \otimes I_2^{\otimes(m-k)} \Big\|_{B(\ell_{2^m}^2)} = \bigg(\sum_{k=1}^m |\alpha_k|^2 \bigg)^{\frac{1}{2}}. \tag{5.4}$$

Let $\mathcal{H} = \ell^2(H) \overset{2}{\oplus} \ell^2(H)$, let $\sigma \colon \ell^2(H) \to \ell^2(H)$ denote the shift operator, let $\Gamma \colon \ell^2(H) \to \ell^2(H)$ be the Hankel operator associated to the $B(H)$-valued function F given by

$$F(t) = \sum_{k=1}^\infty \frac{\widetilde{C_k}}{2^k} \, e^{-i(2^k-1)t},$$

and let $T \in B(\mathcal{H})$ be the operator given by

$$T = \begin{bmatrix} \sigma^* & \Gamma \\ 0 & \sigma \end{bmatrix}.$$

Pisier proved that this operator is polynomially bounded without being completely polynomially bounded. Since $\| \ \|_2 \leq \| \ \|_p$ on $\mathcal{P}$, the linear mapping

$$u \colon (\mathcal{P}, \| \ \|_p) \longrightarrow B(\mathcal{H}), \qquad u(\varphi) = \varphi(T),$$

is therefore bounded. Our aim is now to show that u is not p-completely bounded.

We consider the auxiliary mapping $w \colon \mathcal{P} \to B(H)$ defined by letting

$$w\bigg(\sum_{k \geq 0} d_k z^k \bigg) \ = \ \sum_{k \geq 0} d_{2^k} \widetilde{C_k}$$

for any finite sequence $(d_k)_{k \geq 0}$ of complex numbers.

Let $j \colon H \to \ell^2(H)$ be the isometric embedding given by $j(x) = (x, 0, \ldots, 0, \ldots)$. Then let $v \colon B(\ell^2(H)) \to B(H)$ be defined by letting $v(R) = j^* R j$ for any $R \in B(H)$. It is easy to

check that v is p-completely bounded, with $\|v\|_{pcb} = 1$. On the other hand, for any $\varphi \in \mathcal{P}$, we have

$$\varphi(T) = \begin{bmatrix} \varphi(\sigma^*) & \Gamma\varphi'(\sigma) \\ 0 & \varphi(\sigma) \end{bmatrix},$$

see [47, (9.7)]. Let $\widetilde{u}\colon \mathcal{P} \to B(\ell^2(H))$ be defined by $\widetilde{u}(\varphi) = \Gamma\varphi'(\sigma)$. Then the argument in the proof of [47, Thm. 9.7] shows that $w = v\widetilde{u}$. Thus if u were p-completely bounded, then w would be p-completely bounded as well. Let us show that this does not hold true.

Note that for any Banach space X, for any integer $N \geq 1$, for any $T \in B(X)$ and for any $A \in B(\ell^1_N)$, we have

$$\big\|A \otimes T\colon \ell^1_N(X) \longrightarrow \ell^1_N(X)\big\| \;=\; \|A\|_{B(\ell^1_N)}\|T\|_{B(X)}.$$

This can be be seen as a consequence of the fact that $\ell^1_N(X)$ is the projective tensor product of ℓ^1_N and X, see [15, Chapter VIII], however an elementary proof is also possible (we leave this to the reader).

Let $m \geq 1$. Clearly $\|E\|_{B(\ell^1_2)} = \|D\|_{B(\ell^1_2)} = 1$. Hence applying the above property we have $\|C_k\|_{B(\ell^1_{2^k})} = \|E\|^{k-1}_{B(\ell^1_2)}\|D\|_{B(\ell^1_2)} = 1$ and hence

$$\big\|C_k \otimes I_2^{\otimes(m-k)} \otimes S^{2^k}\big\|_{B(\ell^1_{2^m}(\ell^1_{\mathbb{Z}}))} = 1, \qquad k = 1, \ldots, n. \tag{5.5}$$

Let $\varphi_m \in M_{2^m} \otimes \mathcal{P}$ be given by

$$\varphi_m(z) = \sum_{k=1}^m C_k \otimes I_2^{\otimes(m-k)}\, z^{2^k}.$$

By (5.4), we have

$$\|\varphi_m\|_2 = \sup_{|z|=1} \|\varphi_m(z)\|_{B(\ell^2_{2^m})} = \sup_{|z|=1} \left\| \sum_{k=1}^m z^{2^k}\, C_k \otimes I_2^{\otimes(m-k)} \right\|_{B(\ell^2_{2^m})} = \sqrt{m}.$$

On the other hand, applying (5.5) we have

$$\|\varphi_m\|_1 \;=\; \left\|\sum_{k=1}^m C_k \otimes I_2^{\otimes(m-k)} \otimes S^{2^k}\right\|_{B(\ell^1_{2^m}(\ell^1_{\mathbb{Z}}))} \leq \sum_{k=1}^m \big\|C_k \otimes I_2^{\otimes(m-k)} \otimes S^{2^k}\big\|_{B(\ell^1_{2^m}(\ell^1_{\mathbb{Z}}))} \;=\; m.$$

By interpolation, we deduce that

$$\|\varphi_m\|_p \leq m^{\frac{1}{p}}. \tag{5.6}$$

Next we have

$$\big(I_{B(\ell^p_{2^m})} \otimes w\big)(\varphi_m) \;=\; \sum_{k=1}^m C_k \otimes I_2^{\otimes(m-k)} \otimes \widetilde{C_k}\,. \tag{5.7}$$

Let us estimate the norm of this tensor product in $B(\ell^p_{2^m}(H))$. Let (e_1, e_2) denote the canonical basis of $\mathbb{C}^2$. For any $k = 1, \dots, m$ and any $i_1, \dots, i_m, j_1, \dots, j_m$ in $\{1, 2\}$,

$$\begin{aligned}
&\Big\langle \Big(C_k \otimes I_2^{\otimes(m-k)}\Big)(e_{j_1} \otimes \cdots \otimes e_{j_m}), e_{i_1} \otimes \cdots \otimes e_{i_m} \Big\rangle \\
&\quad = \big\langle Ee_{j_1} \otimes \cdots \otimes Ee_{j_{k-1}} \otimes De_{j_k} \otimes e_{j_{k+1}} \cdots \otimes e_{j_m}, e_{i_1} \otimes \cdots \otimes e_{i_m} \big\rangle \\
&\quad = \big\langle (-1)^{\delta_{j_1,2}} e_{j_1} \otimes \cdots \otimes (-1)^{\delta_{j_{k-1},2}} e_{j_{k-1}} \otimes \delta_{j_k,2} e_1 \otimes e_{j_{k+1}} \cdots \otimes e_{j_m}, e_{i_1} \otimes \cdots \otimes e_{i_m} \big\rangle \\
&\quad = (-1)^{\delta_{j_1,2}} \cdots (-1)^{\delta_{j_{k-1},2}} \delta_{j_k,2} \big\langle e_{j_1}, e_{i_1} \big\rangle \cdots \big\langle e_{j_{k-1}}, e_{i_{k-1}} \big\rangle \big\langle e_1, e_{i_k} \big\rangle \big\langle e_{j_{k+1}}, e_{i_{k+1}} \big\rangle \cdots \big\langle e_{j_m}, e_{i_m} \big\rangle \\
&\quad = (-1)^{\delta_{j_1,2}} \cdots (-1)^{\delta_{j_{k-1},2}} \delta_{j_k,2} \delta_{1,i_k} \delta_{j_1,i_1} \cdots \delta_{j_{k-1},i_{k-1}} \delta_{j_{k+1},i_{k+1}} \cdots \delta_{j_m,i_m}.
\end{aligned}$$

Hence

$$\Big\langle \Big(\sum_{k=1}^m \Big(C_k \otimes I_2^{\otimes(m-k)}\Big) \otimes \Big(C_k \otimes I_2^{\otimes(m-k)}\Big) \Big) \Big(\sum_{j_1,\dots,j_m=1}^2 e_{j_1} \otimes \cdots \otimes e_{j_m} \otimes e_{j_1} \otimes \cdots \otimes e_{j_m} \Big), \sum_{i_1,\dots,i_m=1}^2 e_{i_1} \otimes \cdots \otimes e_{i_m} \otimes e_{i_1} \otimes \cdots \otimes e_{i_m} \Big\rangle$$

is equal to

$$\begin{aligned}
&\sum_{k=1}^m \sum_{\substack{j_1,\dots,j_m,\\ i_1,\dots,i_m=1}}^2 \Big\langle \Big(C_k \otimes I_2^{\otimes(m-k)}\Big)(e_{j_1} \otimes \cdots \otimes e_{j_m}), e_{i_1} \otimes \cdots \otimes e_{i_m} \Big\rangle^2 \\
=&\sum_{k=1}^m \sum_{\substack{j_1,\dots,j_m,\\ i_1,\dots,i_m=1}}^2 \big(\delta_{j_k,2} \delta_{1,i_k} \delta_{j_1,i_1} \cdots \delta_{j_{k-1},i_{k-1}} \delta_{j_{k+1},i_{k+1}} \cdots \delta_{j_m,i_m}\big)^2 \\
=&\sum_{k=1}^m 2^{m-1} \;=\; m 2^{m-1}.
\end{aligned}$$

Since the norm of

$$\sum_{i_1,\dots,i_m=1}^2 e_{i_1} \otimes \cdots \otimes e_{i_m} \otimes e_{i_1} \otimes \cdots \otimes e_{i_m}$$

in $\ell^p_{2^m}(\ell^2_{2^m})$ (resp. in $\ell^{p'}_{2^m}(\ell^2_{2^m})$) is equal to

$$\Big(\sum_{i_1,\dots,i_m=1}^2 \|e_{i_1} \otimes \cdots \otimes e_{i_m}\|^p_{\ell^2_{2^m}} \Big)^{\frac{1}{p}} \;=\; 2^{\frac{m}{p}}$$

(resp. $2^{\frac{m}{p'}}$), we deduce that

$$\Big\| \sum_{k=1}^m \Big(C_k \otimes I_2^{\otimes(m-k)}\Big) \otimes \Big(C_k \otimes I_2^{\otimes(m-k)}\Big) \Big\|_{B(\ell^p_{2^m}(\ell^2_{2^m}))} \geq \frac{m}{2}.$$

Combining with (5.3) and (5.7) we obtain that

$$\big\| \big(I_{B(\ell^p_{2^m})} \otimes w\big)(\varphi_m) \big\| \;\geq\; \frac{m}{2}\,.$$

Together with (5.6), this implies that w is not p-completely bounded. Thus T is not p-completely polynomially bounded.

So we are done except that T acts on the Hilbert space $\mathcal{H}$ and not on $L^p\big([0,1]\big)$. However arguing as in the last part of the proof of Theorem 4.8, it is easy to pass from $\mathcal{H}$ to the space $L^p\big([0,1]\big)$. □

The proof of Proposition 4.6 actually yields the following stronger result: if an operator $T\colon L^p(\Omega) \to L^p(\Omega)$ admits a loose dilation, then it is p-completely polynomially bounded (details are left to the reader). Hence the above proposition yields the following.

Corollary 5.3. *There exists a p-polynomially bounded operator $T\colon L^p\big([0,1]\big) \to L^p\big([0,1]\big)$ which does not admit any loose dilation.*

Note also that according to Theorem 4.8 and the above observation, no R-Ritt operator can satisfy Proposition 5.2. Namely, if $T\colon L^p(\Omega) \to L^p(\Omega)$ is an R-Ritt operator and is p-polynomially bounded, then it is p-completely polynomially bounded.

Remark 4.3 and the above investigations lead to the following open problem (for $p \neq 2$): *does any p-completely polynomially bounded operator $T\colon L^p(\Omega) \to L^p(\Omega)$ admit a loose dilation?*

In the last part of this section we are going to consider another type of counterexamples. Clearly any p-polynomially bounded $T\colon L^p(\Omega) \to L^p(\Omega)$ is automatically power bounded, that is,

$$\sup_{n\geq 0} \|T^n\| \,<\infty\,.$$

The existence of a power bounded operator on Hilbert space which is not polynomially bounded is an old result of Foguel [18]. Our aim is to prove an L^p-analog of that result. We will actually show a stronger form: there exists a Ritt operator which is not polynomialy bounded. To achieve this, we will adapt a technique from [24].

We need some background on Schauder bases and their multipliers that we briefly recall. We let v_1 denote the set of all sequences $(c_n)_{n\geq 0}$ of complex numbers whose variation $\sum_{n=1}^{\infty} |c_n - c_{n-1}|$ is finite. Any such sequence is bounded and v_1 is a Banach space for the norm

$$\big\|(c_n)_{n\geq 0}\big\|_{v_1} \;=\; |c_0| \,+\, \sum_{n=1}^{\infty} |c_n - c_{n-1}|\,.$$

Let $(e_n)_{n\geq 0}$ be a Schauder basis on some Banach space X. For any $n \geq 0$, let $Q_n\colon X \to X$ be the projection defined by

$$Q_n\Big(\sum_{k=0}^{\infty} a_k e_k\Big) \;=\; \sum_{k=n}^{\infty} a_k e_k \tag{5.8}$$

for any converging sequence $\sum_k a_k e_k$. The sequence $(Q_n)_{n\geq 0}$ is bounded and by a standard Abel summation argument, we have the following.

Lemma 5.4. *For any $c=(c_n)_{n\geq 0}$ in v_1, there exists a (necessarily unique) bounded operator $T_c\colon X\to X$ such that*

$$T_c\Big(\sum_{n=0}^{\infty} a_n e_n\Big) \,=\, \sum_{n=0}^{\infty} c_n a_n e_n$$

for any converging sequence $\sum_n a_n e_n$. Furthermore,

$$\|T_c\| \leq \Big(\sup_{n\geq 0}\|Q_n\|\Big)\big\|(c_n)_{n\geq 1}\big\|_{v_1}.$$

The above operator T_c is called the multiplier associated to the sequence c.

Proposition 5.5. *Let $1<p<\infty$.*

1. *There exists a Ritt (hence a power bounded) operator on ℓ^2 which is not p-polynomially bounded.*
2. *There exists an R-Ritt operator on $L^p\big([0,1]\big)$ which is not p-polynomially bounded.*

Proof. (1): We let $(e_n)_{n\geq 0}$ be a Schauder basis of $H=\ell^2$. It is clear that the sequence $\big(1-\frac{1}{2^n}\big)_{n\geq 0}$ has a finite variation. According to the above discussion, we let $T\colon H\to H$ denote the multiplier associated to that sequence.

For any $\theta\in(-\pi,0)\cup(0,\pi]$, set

$$c(\theta)_n \,=\, \frac{1}{e^{i\theta}-\big(1-\frac{1}{2^n}\big)}\,, \qquad n\geq 0.$$

We have

$$\begin{aligned}\sum_{n=1}^{+\infty}|c(\theta)_n-c(\theta)_{n-1}| \,&=\, \sum_{n=1}^{+\infty}\Big|\int_{1-\frac{1}{2^{n-1}}}^{1-\frac{1}{2^n}}\frac{dt}{\big(e^{i\theta}-t\big)^2}\Big|\\ &\leq \sum_{n=1}^{+\infty}\int_{1-\frac{1}{2^{n-1}}}^{1-\frac{1}{2^n}}\frac{dt}{|e^{i\theta}-t|^2}\\ &\leq \int_0^1\frac{dt}{|e^{i\theta}-t|^2}\,.\end{aligned}$$

Let $I(\theta)$ denote the latter integral. It is finite hence $c(\theta)=(c(\theta)_n)_{n\geq 0}$ belongs to v_1. It is easy to deduce that $e^{i\theta}-T$ is invertible, the operator $R(e^{i\theta},T)$ being the multiplier associated to the sequence $c(\theta)$.

For $\theta\neq\pi$, elementary computations yield

$$\begin{aligned}I(\theta) \,&=\, \int_0^1\frac{dt}{(t-\cos(\theta))^2+\sin^2(\theta)}\\ &=\, \frac{1}{\sin(\theta)}\int_{-\frac{\cos(\theta)}{\sin(\theta)}}^{\frac{1-\cos(\theta)}{\sin(\theta)}}\frac{du}{1+u^2}\\ &=\, \frac{\pi-\theta}{2\sin(\theta)}\,.\end{aligned}$$

Moreover $|e^{i\theta}-1| = 2\sin\big(\frac{\theta}{2}\big)$, hence

$$|e^{i\theta}-1|I(\theta) \;=\; (\pi-\theta)\frac{\sin\big(\frac{\theta}{2}\big)}{\sin(\theta)} \;=\; \frac{\pi-\theta}{2\cos\big(\frac{\theta}{2}\big)}\,.$$

This is bounded for θ varying in $(-\pi,0)\cup(0,\pi)$. According to Lemma 5.4, this shows that

$$\sigma(T)\subset\mathbb{D}\cup\{1\} \qquad \text{and} \qquad \big\{(\lambda-1)R(\lambda,T)\,:\,\lambda\in\mathbb{T}\setminus\{1\}\big\} \quad \text{is bounded.}$$

Applying the maximum principle to the function $z\mapsto(1-z)(I_H-zT)^{-1}$, we deduce that the set $\{(\lambda-1)R(\lambda,T)\,:\,|\lambda|>1\}$ is bounded as well, and hence T is a Ritt operator.

Let us now assume that the basis $(e_n)_{n\geq 0}$ is not an unconditional one. The operator $I-T$ is the multiplier associated to the sequence $\big(\frac{1}{2^n}\big)_{n\geq 0}$ and as is well-known, the lack of unconditionality implies that for any $\theta\in(0,\pi)$, this operator does not have a bounded $H^\infty(\Sigma_\theta)$ functional calculus (see e.g. [32, Thm. 4.1] and its proof). According to Proposition 4.7, this implies that T is not p-polynomially bounded.

(2): Since all bounded subsets of $B(\ell^2)$ are R-bounded, the operator considered in part (1) is automatically an R-Ritt operator. Then arguing again as in the proof of Theorem 4.8, it is easy to pass from an ℓ^2-operator to an $L^p\big([0,1]\big)$-operator which is not p-polynomially bounded although being an R-Ritt operator. □

6. Generalizations to general Banach spaces

Up to now we have mostly dealt with operators acting on (commutative) L^p-spaces. In this last section, we shall consider more general Banach spaces, in particular noncommutative L^p-spaces. We aim at extending our main results from Sections 3 and 4 to this broader context.

We will use classical notions from Banach space theory such as cotype, K-convexity and the UMD property. We refer the reader to [8, 14, 43] for background.

In accordance with (2.1), we are going to extend the definitions (1.1) and (1.3) to arbitrary Banach spaces using Rademacher averages. Recall Section 2 for notation. The use of such averages as a substitute of square functions on abstract Banach spaces is a classical and fruitful principle. See e.g. [22, 23, 30].

Let X be a Banach space, let $T\colon X\to X$ be any bounded operator and let $x\in X$. Consider the element $x_k = k^{\frac{1}{2}}(T^k(x)-T^{k-1}(x))$ for any $k\geq 1$. If the series $\sum_k \varepsilon_k\otimes x_k$ converges in $L^2(\Omega_0;X)$ then we set

$$\|x\|_{T,1} \;=\; \Big\|\sum_{k=1}^{\infty} k^{\frac{1}{2}}\,\varepsilon_k\otimes\big(T^k(x)-T^{k-1}(x)\big)\Big\|_{\mathrm{Rad}(X)}\,.$$

We set $\|x\|_{T,1}=\infty$ otherwise. Likewise, if T is a Ritt operator and $\alpha>0$ is a positive real number, then we set

$$\|x\|_{T,\alpha} \;=\; \Big\|\sum_{k=1}^{\infty} k^{\alpha-\frac{1}{2}}\,\varepsilon_k\otimes T^{k-1}(I-T)^\alpha x\Big\|_{\mathrm{Rad}(X)}$$

if the corresponding series converges in $L^2(\Omega_0;X)$, and $\|x\|_{T,\alpha}=\infty$ otherwise. The following extends Theorem 3.3.

Theorem 6.1. *Assume that X is reflexive and has a finite cotype. Let $T\colon X\to X$ be an R-Ritt operator. Then for any $\alpha>0$ and $\beta>0$, we have an equivalence*

$$\|x\|_{T,\alpha}\approx\|x\|_{T,\beta},\qquad x\in X.$$

Proof. We noticed in Section 2 that if X has a finite cotype, then Rademacher averages and Gaussian averages are equivalent on X.

Furthermore, the reflexivity of X ensures that it satisfies the Mean Ergodic Theorem. We thus have

$$X=\operatorname{Ker}(I-T)\oplus\overline{\operatorname{Ran}(I-T)}.$$

Lastly, since X has a finite cotype, it cannot contain c_0 (as an isomorphic subspace). Hence by [27], a series $\sum_k \varepsilon_k\otimes x_k$ converges in $L^2(\Omega_0;X)$ if (and only if) its partial sums are uniformly bounded, that is, there is a constant $K\geq 0$ such that

$$\Big\|\sum_{k=1}^N \varepsilon_k\otimes x_k\Big\|_{\mathrm{Rad}(X)}\leq K,\qquad N\geq 1.$$

With these three properties in hand, it is easy to see that our proof of Theorem 3.3 extends verbatim to the general case. □

In the rest of this section we are going to focus on noncommutative L^p-spaces. We let M be a semifinite von Neumann algebra equipped with a normal semifinite faithful trace and for any $1\leq p<\infty$, we let $L^p(M)$ denote the associated (noncommutative) L^p-space. We refer to [49] for background and information on these spaces. Any element of $L^p(M)$ is a (possibly unbounded) operator and for any such x, we set

$$|x|=(x^*x)^{\frac12}.$$

We recall the noncommutative analog of (2.1) from [28] (see also [29]). For finite families $(x_k)_k$ of $L^p(M)$, we have the following equivalences. If $2\leq p<\infty$, then

$$\Big\|\sum_k \varepsilon_k\otimes x_k\Big\|_{\mathrm{Rad}(L^p(M))}\approx\max\Big\{\Big\|\Big(\sum_k|x_k|^2\Big)^{\frac12}\Big\|_{L^p(M)},\Big\|\Big(\sum_k|x_k^*|^2\Big)^{\frac12}\Big\|_{L^p(M)}\Big\}. \tag{6.1}$$

If $1<p\leq 2$, then

$$\Big\|\sum_k \varepsilon_k\otimes x_k\Big\|_{\mathrm{Rad}(L^p(M))}\approx\inf\Big\{\Big\|\Big(\sum_k|u_k|^2\Big)^{\frac12}\Big\|_{L^p(M)}+\Big\|\Big(\sum_k|v_k^*|^2\Big)^{\frac12}\Big\|_{L^p(M)}\Big\}, \tag{6.2}$$

where the infimum runs over all possible decompositions $x_k=u_k+v_k$ in $L^p(M)$.

Let $T\colon L^p(M)\to L^p(M)$ be a bounded operator. We say that T admits a noncommutative loose dilation if there exist a von Neumann algebra $\widetilde{M}$, an isomorphism $U\colon L^p(\widetilde{M})\to L^p(\widetilde{M})$ such that the set $\{U^n\,:\,n\in\mathbb{Z}\}$ is bounded and two bounded maps $L^p(M)\xrightarrow{J}L^p(\widetilde{M})$ and $L^p(\widetilde{M})\xrightarrow{Q}L^p(M)$ such that $T^n=QU^nJ$ for any integer $n\geq 0$. We say that T admits a noncommutative strict dilation if this holds true for an isometric isomorphism U and two

contractions J and Q. As opposed to the commutative case (see Remark 4.2), there is no characterization of contractions $T\colon L^p(M) \to L^p(M)$ which admit a noncommutative strict dilation. The gap with the commutative situation is illustrated by the following result [21, Thm 5.1]: for any $p \neq 2$, there exist a completely positive contraction on some finite dimensional noncommutative L^p which does not admit any noncommutative strict dilation.

We now turn to loose dilations. In the commutative setting, the following proposition is a combination of Propositions 4.6 and 4.7.

Proposition 6.2. *Let $T\colon L^p(M) \to L^p(M)$, with $1 < p < \infty$. If T admits a noncommutative loose dilation, then $I - T$ is sectorial and admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus for any $\theta \in \big(\frac{\pi}{2}, \pi\big)$.*

Proof. Let us explain how to adapt the 'commutative' proof to the present setting.

First we extend the definition (4.1) as follows. For any Banach space X, let $S_X\colon \ell^p_{\mathbb{Z}}(X) \to \ell^p_{\mathbb{Z}}(X)$ denote the shift operator. Then for any $\varphi \in \mathcal{P}$, we set

$$\|\varphi\|_{p,X} = \big\|\varphi(S_X)\big\|_{B(\ell^p_{\mathbb{Z}}(X))}.$$

It follows from [6, Thm. 4.3] that if X is UMD, then Theorem 4.4 holds as well for scalar valued Fourier multipliers on $\ell^p_{\mathbb{Z}}(X)$. In this case, the argument in the proof of Proposition 4.7 leads to the following: for any $\theta \in \big(\frac{\pi}{2}, \pi\big)$, there is an estimate

$$\|\varphi\|_{p,X} \lesssim \|\varphi\|_{H^\infty(\mathbb{D}_\theta)} \tag{6.3}$$

for rational functions φ with poles outside $\overline{\mathbb{D}_\theta}$.

Second we note that if $U\colon L^p(\widetilde{M}) \to L^p(\widetilde{M})$ is an isomorphism such that $K = \sup\{\|U^n\| : n \in \mathbb{Z}\} < \infty$, then the vectorial version of the transference principle (see [4, Thm. 2.8]) ensures that for any φ as above, we have

$$\|\varphi(U)\| \leq K^2 \|\varphi\|_{p, L^p(\widetilde{M})}.$$

Assume now that $T\colon L^p(M) \to L^p(M)$ admits a noncommutative loose dilation. Noncommutative L^p-spaces are UMD hence property (6.3) applies to them. Hence arguing as in proof of Proposition 4.6, we find an estimate

$$\|\varphi(T)\| \lesssim \|\varphi\|_{H^\infty(\mathbb{D}_\theta)}$$

for rational functions φ with poles outside $\overline{\mathbb{D}_\theta}$. Finally the argument at the end of the proof of Proposition 4.7 yields that $I - T$ admits a bounded $H^\infty(\Sigma_\theta)$ functional calculus for any $\theta > \frac{\pi}{2}$. We skip the details. □

We are now ready to give the noncommutative analog of Theorem 4.8.

Theorem 6.3. *Let $T\colon L^p(M) \to L^p(M)$ be a bounded operator, with $1 < p < \infty$.*

(1) *The following assertions are equivalent.*
   (i) *The operator T is R-Ritt and admits a noncommutative loose dilation.*
   (ii) *The operator T and its adjoint $T^*\colon L^{p'}(M) \to L^{p'}(M)$ both satisfy uniform estimates*

$$\|x\|_{T,1} \lesssim \|x\|_{L^p(M)} \qquad \text{and} \qquad \|y\|_{T^*,1} \lesssim \|y\|_{L^{p'}(M)}$$

   for $x \in L^p(M)$ and $y \in L^{p'}(M)$.

(2) *Assume that $p \geq 2$. Then the above conditions are equivalent to the existence of a constant $C \geq 1$ for which the following two properties hold.*

(iii) *For any $x \in L^p(M)$,*

$$\Big\| \Big(\sum_{k=1}^{\infty} k \big| T^k(x) - T^{k-1}(x) \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p(M)} \leq C \|x\|_{L^p(M)}$$

and

$$\Big\| \Big(\sum_{k=1}^{\infty} k \big| \big(T^k(x) - T^{k-1}(x)\big)^* \big|^2 \Big)^{\frac{1}{2}} \Big\|_{L^p(M)} \leq C \|x\|_{L^p(M)}.$$

(iii)* *For any $y \in L^{p'}(M)$, there exist two sequences $(u_k)_{k\geq 1}$ and $(v_k)_{k\geq 1}$ of $L^{p'}(M)$ such that*

$$\Big\| \Big(\sum_{k=1}^{\infty} |u_k|^2 \Big)^{\frac{1}{2}} \Big\|_{L^{p'}(M)} \leq C \|y\|_{L^{p'}(M)}\,, \qquad \Big\| \Big(\sum_{k=1}^{\infty} |v_k^*|^2 \Big)^{\frac{1}{2}} \Big\|_{L^{p'}(M)} \leq C \|y\|_{L^{p'}(M)}\,,$$

and

$$u_k + v_k \,=\, k^{\frac{1}{2}} \big(T^{*k}(y) - T^{*(k-1)}(y) \big) \qquad \text{for any } k \geq 1.$$

Proof. Theorem 3.1 holds as well on noncommutative L^p-spaces, by [30]. Combining that result with Proposition 6.2, we obtain that (i) implies (ii).

Assume (ii) and suppose for simplicity that $I - T$ is 1-1 (the changes to treat the general case are minor ones). By Theorem 6.1, we have uniform estimates

$$\|x\|_{T,\frac{1}{2}} \,\lesssim\, \|x\|_{L^p(M)} \qquad \text{and} \qquad \|y\|_{T^*,\frac{1}{2}} \,\lesssim\, \|y\|_{L^{p'}(M)}$$

for $x \in L^p(M)$ and $y \in L^{p'}(M)$. As in the proof of Theorem 4.8, we may therefore define $J_1 \colon L^p(M) \to \mathrm{Rad}(L^p(M))$ and $J_2 \colon L^{p'}(M) \to \mathrm{Rad}(L^{p'}(M))$ by setting

$$J_1(x) = \sum_{k=1}^{\infty} \varepsilon_k \otimes T^{k-1}(I-T)^{\frac{1}{2}} x \qquad \text{and} \qquad J_2(y) = \sum_{k=1}^{\infty} \varepsilon_k \otimes T^{*(k-1)}(I-T^*)^{\frac{1}{2}} y$$

for any $x \in L^p(M)$ and any $y \in L^{p'}(M)$. Since $L^p(M)$ is K-convex, we have a natural isomorphism

$$\big(\mathrm{Rad}(L^p(M)) \big)^* \,\approx\, \mathrm{Rad}(L^{p'}(M)). \tag{6.4}$$

Hence one can consider the composition $J_2^* J_1$, it is equal to $(I+T)^{-1}$ and one obtains (i) by simply adapting the proof of Theorem 4.8.

Finally the equivalence between (ii) and (iii)+(iii)* follows from (6.1) and (6.2). □

Switching (iii) and (iii)*, we find a version of (2) for the case $p \leq 2$.

Remark 6.4.

(1) Let $T \colon L^p(\Omega) \to L^p(\Omega)$ be an R-Ritt operator on some *commutative* L^p-space. Combining Theorems 6.3 and 4.8, we find that T admits a noncommutative loose dilation (if and) only if it admits a commutative one.

(2) Proposition 3.4 holds true on noncommutative L^p-spaces. The proof is similar, using (6.4) instead of the duality $L^p(\Omega;\ell^2)^* = L^{p'}(\Omega;\ell^2)$.

We refer the reader to the forthcoming paper [3] for more about square functions associated to Ritt operators on noncommutative L^p-spaces.

References

[1] M. Akcoglu, *A pointwise ergodic theorem in L_p-spaces*, Canad. J. Math. 27 (1975), no. 5, 1075-1082.

[2] M. Akcoglu, and L. Sucheston, *Dilations of positive contractions on L_p spaces*, Canad. Math. Bull. 20 (1977), 285-292.

[3] C. Arhancet, *Square functions for Ritt operators on noncommutative L^p-spaces*, Preprint 2011.

[4] E. Berkson, and T. A. Gillespie, *Generalized analyticity in UMD spaces*, Ark. Mat. 27 (1989), 1-14.

[5] E. Berkson, and T. A. Gillespie, *Spectral decompositions and harmonic analysis on UMD Banach spaces*, Studia Math. 112 (1994), 13-49.

[6] S. Blunck, *Maximal regularity of discrete and continuous time evolution equations*, Studia Math. 146 (2001), no. 2, 157-176.

[7] S. Blunck, *Analyticity and discrete maximal regularity on L_p-spaces*, J. Funct. Anal. 183 (2001), 211-230.

[8] D. L. Burkholder, *Martingales and singular integrals in Banach spaces*, pp. 233-269 in "Handbook of the geometry of Banach spaces", Vol. I, North-Holland, Amsterdam, 2001.

[9] P. Clément, B. de Pagter, F. A. Sukochev, and H. Witvliet, *Schauder decompositions and multiplier theorems*, Studia Math. 138 (2000), 135-163.

[10] R. Coifman, R. Rochberg, and G. Weiss, *Applications of transference: the L_p version of von Neumann's inequality and the Littlewood-Paley-Stein theory*, pp. 53-67 in "Linear spaces and Approximation", Birkhäuser, Basel, 1978.

[11] R. R. Coifman, and G. Weiss, *Transference methods in analysis*, CBMS 31, Amer. Math. Soc., 1977.

[12] M. Cowling, I. Doust, A. McIntosh, and A. Yagi, *Banach space operators with a bounded H^∞ functional calculus*, J. Aust. Math. Soc., Ser. A 60 (1996), 51-89.

[13] K. Davidson, *C^*-algebras by example*, Fields Institute Monographs, 6. American Mathematical Society, Providence, RI, 1996. xiv+309 pp.

[14] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely summing operators*, Cambridge Studies in Advanced Mathematics 43, Cambridge University Press, Cambridge, 1995. xvi+474 pp.

[15] J. Diestel, and J. J. Uhl, *Vector measures*, Mathematical Surveys, No. 15, American Mathematical Society, Providence, R.I., 1977. xiii+322 pp.

[16] S. W. Drury, *A counterexample to a conjecture of Matsaev*, Lin. Alg. and its Appl. 435 (2011), 323-329.

[17] R. E. Edwards, and G. I. Gaudry, *Littlewood-Paley and multiplier theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 90. Springer-Verlag, Berlin-New York, 1977. ix+212 pp.

[18] S. R. Foguel, *A counterexample to a problem of Sz.-Nagy*, Proc. Amer. Math. Soc. 15 (1964), 788-790.

[19] A. M. Fröhlich, and L. Weis, *H^∞ calculus and dilations*, Bull. Soc. Math. France 134 (2006), no. 4, 487-508.

[20] M. Haase, *The functional calculus for sectorial operators*, Operator Theory: Advances and Applications, 169, Birkhuser Verlag, Basel, 2006. xiv+392 pp.

[21] M. Junge, and C. Le Merdy, *Dilations and rigid factorisations on noncommutative L^p-spaces*, J. Funct. Anal. 249 (2007), no. 1, 220-252.

[22] M. Junge, C. Le Merdy, and Q. Xu, *H^∞ functional calculus and square functions on noncommutative L^p-spaces*, Soc. Math. France, Astérisque 305, 2006.

[23] N. J. Kalton, Nigel, P. Kunstmann, and L. Weis, *Perturbation and interpolation theorems for the H^∞-calculus with applications to differential operators*, Math. Ann. 336 (2006), no. 4, 747-801.

[24] N. J. Kalton, and C. Le Merdy, *Solution of a problem of Peller concerning similarity*, J. Operator Theory 47 (2002), no. 2, 379-387.

[25] U. Krengel, *Ergodic theorems*, de Gruyter Studies in Mathematics, 6. Walter de Gruyter and Co., Berlin, 1985. viii+357 pp.

[26] P. C. Kunstmann, and L. Weis, *Maximal L_p-regularity for parabolic equations, Fourier multiplier theorems and H^∞-functional calculus*, pp. 65-311 in "Functional analytic methods for evolution equations", Lect. Notes in Math. 1855, Springer, 2004.
[27] S. Kwapien, *On Banach spaces containing c_0*, Studia Math. 52 (1974), 187-188.
[28] F. Lust-Piquard, *Inégalités de Khintchine dans C_p $(1 < p < \infty)$* (French), C. R. Acad. Sci. Paris Sr. I Math. 303 (1986), no. 7, 289-292.
[29] F. Lust-Piquard, and G. Pisier, *Noncommutative Khintchine and Paley inequalities*, Ark. Mat. 29 (1991), no. 2, 241-260.
[30] C. Le Merdy, *H^∞ functional calculus and square function estimates for Ritt operators*, Preprint 2011.
[31] C. Le Merdy, *Square functions, bounded analytic semigroups, and applications*, Banach Center Publ. 75 (2007), 191-220.
[32] C. Le Merdy, *H^∞-functional calculus and applications to maximal regularity*, Publ. Math. Besançon 16 (1998), 41-77.
[33] C. Le Merdy, *On square functions associated to sectorial operators*, Bull. Soc. Math. France 132 (2004), no. 1, 137-156.
[34] C. Le Merdy, and Q. Xu, *Maximal theorems and square functions for analytic operators on L^p-spaces*, Preprint 2010, arXiv:1011.1360.
[35] C. Le Merdy, and Q. Xu, *Strong q-variation inequalities for analytic semigroups*, Preprint 2010, arXiv:1103.2874.
[36] Yu. Lyubich, *Spectral localization, power boundedness and invariant subspaces under Ritt's type condition*, Studia Math. 134 (1999), 153-167.
[37] A. McIntosh, *Operators which have an H^∞ functional calculus*, Proc. CMA Canberra 14 (1986), 210-231.
[38] B. Nagy, and J. Zemanek, *A resolvent condition implying power boundedness*, Studia Math. 134 (1999), 143-151.
[39] O. Nevanlinna, *Convergence of iterations for linear equations*, Birkhaüser, Basel, 1993.
[40] V. I. Paulsen, *Every completely polynomially bounded operator is similar to a contraction*, J. Funct. Anal. 55 (1984), no. 1, 1-17.
[41] V. I. Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics, 78, Cambridge University Press, Cambridge, 2002. xii+300 pp.
[42] V. Peller, *An analogue of J. von Neumann's inequality for the space L^p* (Russian), Dokl. Akad. Nauk SSSR 231 (1976), no. 3, 539-542.
[43] G. Pisier, *Probabilistic methods in the geometry of Banach spaces*, pp. 167241 in "Probability and analysis (Varenna, 1985)", Lecture Notes in Math. 1206, Springer, Berlin, 1986.
[44] G. Pisier, *Complex interpolation and regular operators between Banach lattices*, Arch. Math. (Basel) 62 (1994), no. 3, 2611-269.
[45] G. Pisier, *Completely bounded maps between sets of Banach space operators*, Indiana Univ. Math. J. 39 (1990), no. 1, 249-277.
[46] G. Pisier, *A polynomially bounded operator on Hilbert space which is not similar to a contraction*, J. Amer. Math. Soc. 10 (1997), no. 2, 351-369.
[47] G. Pisier, *Similarity problems and completely bounded maps* (Second, expanded edition), Lecture Notes in Mathematics, 1618. Springer-Verlag, Berlin, 1996. viii+156 pp.
[48] G. Pisier, *Complex Interpolation between Hilbert, Banach and Operator spaces*, Mem. Amer. Math. Soc. 208 (2010) vi+78 pp.
[49] G. Pisier, and Q. Xu, *Non-commutative L^p-spaces*, pp. 1459-1517 in "Handbook of the Geometry of Banach Spaces", Vol. II, edited by W.B. Johnson and J. Lindenstrauss, Elsevier, 2003.
[50] E.M. Stein, *Topics in harmonic analysis related to the Littlewood-Paley theory*, Ann. Math. Studies, Princeton, University Press, 1970.
[51] B. Sz.-Nagy, and C. Foias, *Harmonic analysis of operators on Hilbert space*, Akademiai Kiadó, Budapest, 1970.
[52] P. Vitse, *A band limited and Besov class functional calculus for Tadmor-Ritt operators*, Archiv. Math. 85 (2005), no. 4, 374-385.

Laboratoire de Mathématiques, Université de Franche-Comté, 25030 Besançon Cedex, France
E-mail address: clemerdy@univ-fcomte.fr

Laboratoire de Mathématiques, Université de Franche-Comté, 25030 Besançon Cedex, France
E-mail address: cedric.arhancet@univ-fcomte.fr